Correlation functions after a quantum quench in a spin-chain in the pre-thermalized regime

Aditi Mitra
Department of Physics, New York University, 4 Washington Place, New York, New York 10003, USA

(Dated: February 14, 2013)

arXiv:1302.2953v1 [cond-mat.str-el] 12 Feb 2013

Results are presented for a two-point correlation function of a spin-chain after a quantum quench for an intermediate time regime where inelastic effects are weak. A Callan-Symanzik like equation for the correlation function is explicitly constructed which is used to show the appearance of two scaling regimes. One is for spatial separations within a light-cone, and the second is for spatial separations on the light cone. In both these regimes, the correlation function is found to decay with power-laws with new nonequilibrium exponents that differ from those in equilibrium, as well as from those obtained from a quadratic Luttinger liquid theory.

PACS numbers: 05.70.Ln,03.75.Kk,37.10.Jk,71.10.Pm

Motivated by experiments in cold atomic gases [1] and ultra-fast spectroscopy of strongly correlated materials [2, 3], the nonequilibrium dynamics of interacting quantum systems has become a topic at the forefront of research. In this context, dynamics arising due to a quantum quench, where a system is prepared in the ground state of an initial Hamiltonian H_i, and then time-evolved with respect to a final Hamiltonian H_f, is of particular interest because of its potential for addressing several fundamental questions [4]. Some of these are, the mechanisms and time-scales for thermalization [5–10], the possibility of an intermediate time prethermalized regime [11–13], dynamical phase transitions associated with non-analytic behavior during the time evolution [14, 15], dynamics of integrable models [16–21] and the possibility of describing their steady-state in terms of a generalized Gibbs ensemble (GGE) [22]. Yet another important question, which is related to the topic of this paper, is the appearance of universal behavior in the dynamics [4, 23, 24], with the possibility of capturing such a behavior in a renormalization group (RG) approach, even though the system is out of equilibrium.

We study a quantum quench in a generic one-dimensional spin-chain $H = \sum_i \left[S_i^x S_{i+1}^x + S_i^y S_{i+1}^y + \Delta S_i^z S_{i+1}^z\right] + n.n.n.$ where *n.n.n.* denote additional next-nearest-neighbor couplings. We study the dynamics in a continuum theory by taking into account the effect of the leading irrelevant operator. In particular we study the time-evolution of the two-point correlation function of the staggered spin component $R(j) = (-1)^j \langle S_0^z S_j^z\rangle$ by explicitly deriving a Callan-Symanzik (CS) like equation for R, thus revealing conditions under which scaling holds in the nonequilibrium problem. We show that near the critical point, where the interaction is marginally irrelevant, logarithmic corrections arise (as in equilibrium [25–27]), however the nature of the logarithmic corrections are qualitatively different for the nonequilibrium problem.

We study a quench where initially the system is in the ground-state of a Luttinger liquid [28]

$$H_i = \frac{u_0}{2\pi}\int dx \left[K_0\left(\pi\Pi(x)\right)^2 + \frac{1}{K_0}\left(\partial_x\phi(x)\right)^2\right] \quad (1)$$

$-\partial_x\phi/\pi$ represents the density, Π is the variable canonically conjugate to ϕ, K_0 is the dimensionless interaction parameter, and u_0 is the velocity of the sound modes. The system is driven out of equilibrium via an interaction quench at t=0 from $K_0 \to K$, with the leading irrelevant operator corresponding to a commensurate lattice V_{sg} also switched on suddenly, at the same time as the quench. This triggers non-trivial time-evolution from $t > 0$ due to $H_f = H_{f0} + V_{sg}$, where $H_{f0} = H_i(K_0 \to K, u_0 \to u)$, $V_{sg} = -\frac{gu}{\alpha^2}\int dx \cos(\gamma\phi)$, and $\Lambda = u/\alpha$ is a short-distance cut-off. We assume that the quench preserves Galilean invariance so that uK=u_0K_0. We study the time-evolution of the equal time two-point correlation function of the staggered spin component,

$$R(x_1t, x_2t) = 4\langle\cos\left(\frac{\gamma\phi(x_1t)}{2}\right)\cos\left(\frac{\gamma\phi(x_2t)}{2}\right)\rangle \quad (2)$$

Denoting $\hat{O}(xt) = 2\cos\left(\frac{\gamma\phi(xt)}{2}\right)$, R may be written as a Keldysh path-integral representing the time-evolution from the initial pure state $|\psi_i\rangle$ (hence an initial density matrix $\rho = |\psi_i\rangle\langle\psi_i|$) corresponding to the ground state of H_i, $R(x_1t, x_2t) = Tr\left[\rho(t)O(x_10)O(x_20)\right]$ which may be written as $R = Tr\left[e^{-iH_ft}|\psi_i\rangle\langle\psi_i|e^{iH_ft}\hat{O}(x_1)\hat{O}(x_2)\right] = \int \mathcal{D}\left[\phi_{cl}, \phi_q\right] e^{i(S_0+S_{sg})}\hat{O}_I(x_1t)\hat{O}_I(x_2t)$, where $\hat{O}_I$ is the operator in the interaction representation of H_{f0}, S_0 describes the nonequilibrium Luttinger liquid (g=0) [29],

$$S_0 = \frac{1}{2}\int_{-\infty}^{\infty}dx_1\int_{-\infty}^{\infty}dx_2\int_0^t dt_1\int_0^t dt_2 \begin{pmatrix}\phi_{cl}(1) & \phi_q(1)\end{pmatrix} \begin{pmatrix} 0 & G_A^{-1}(1,2) \\ G_R^{-1}(1,2) & -\left[G_R^{-1}G_KG_A^{-1}\right](1,2)\end{pmatrix}\begin{pmatrix}\phi_{cl}(2) \\ \phi_q(2)\end{pmatrix} \quad (3)$$

and S_{sg}=$\frac{gu}{\alpha^2}\int_{-\infty}^{\infty}dx_1\int_0^t dt_1\left[\cos(\gamma\phi_-(1)) - \cos(\gamma\phi_+(1))\right]$. Above $1(2)$=$(x_{1(2)}, t_{1(2)})$, $\phi_{cl,q}$=$\frac{\phi_-\pm\phi_+}{\sqrt{2}}$ with $-/+$ representing fields that are time/anti-time ordered on the Keldysh contour [30].

In a global quench like the one we study, the system is translationally invariant in space, so that $R(x_1 t, x_2 t) = R(x_1 - x_2, t)$. We define the following three exponents which play an important role in the dynamics,

$$K_{eq} = \frac{\gamma^2 K}{4}; K_{neq} = \frac{\gamma^2}{8} K_0 \left(1 + \frac{K^2}{K_0^2}\right)$$
$$K_{tr} = \frac{\gamma^2}{8} K_0 \left(1 - \frac{K^2}{K_0^2}\right) \qquad (4)$$

K_{eq} governs the power-law decay in the ground state of H_{f0}, K_{neq} determines the power-law decay at long times after an interaction quench in the Luttinger liquid [8, 31], K_{tr} determines the crossover from a short-time behavior determined primarily by the initial Luttinger parameter $K_{neq} + K_{tr} = \frac{\gamma^2 K_0}{4}$, to the long time behavior determined by K_{neq} [15]. The role played by these exponents is apparent in R which for $g = 0$ and in the scaling limit (setting u=1 and $r = |x_1 - x_2|, t$ are in units of Λ) $r, t \gg 1$ and far from the light cone ($|r \pm t| \gg 1$) is [15, 31],

$$R^{(0)}(r,t) = \left[\left(\frac{1}{r^2}\right)^{\frac{\gamma^2 K_0}{4}} \left(\left|\frac{r^2 - (2t)^2}{r^2 (2t)^2}\right|\right)^{-K_{tr}}\right]^{1/4} \qquad (5)$$

R shows the "horizon-effect" where outside the light-cone $r \gg 2t$, the correlator depends only on the initial wave-function (and hence the initial Luttinger parameter K_0) albeit with a time-dependent prefactor $t^{K_{tr}/2}$. However within the light-cone $r \ll 2t$, the correlator reaches a steady-state characterized by the nonequlibrium exponent K_{neq}. Moreover, on the light-cone $R^{(0)}(r = 2t) \simeq \left[\left(\frac{1}{r^2}\right)^{\frac{\gamma^2 K_0}{4}} \left(\frac{1}{r^3}\right)^{-K_{tr}}\right]^{1/4}$. It is interesting to contrast this behavior with that of a correlator in a quench from an initially gapped state [20, 32]. For the latter the horizon effect is more pronounced as R is exponentially suppressed outside the light-cone. In contrast for our case, where the quench is from an initially gapless state, the change in the behavior of R from outside the light-cone to inside is less dramatic.

Before we discuss the effects of the cosine potential V_{sg} on the correlator, we present results for the β-function [15, 29]

$$\frac{dg}{d\ln l} = g\left[2 - \left(K_{neq} + \frac{K_{tr}}{1 + 4T_m^2}\right)\right] \qquad (6)$$

$$\frac{dK^{-1}}{d\ln l} = \frac{\pi g^2 \gamma^2}{4} I_K(T_m) \qquad (7)$$

$$\frac{d\eta}{d\ln l} = \eta + \frac{\pi g^2 \gamma^2 K}{2} I_\eta(T_m) \qquad (8)$$

$$\frac{d(\eta T_{eff})}{d\ln l} = 2\eta T_{eff} + \frac{\pi g^2 \gamma^2 K}{4} I_{T_{eff}}(T_m) \qquad (9)$$

$$\frac{dT_m}{d\ln l} = -T_m \qquad (10)$$

Eq. (6) shows that at long times after the quench ($T_m \gg 1$), the periodic potential V_{sg} has a nonequilbrium scaling dimension given by K_{neq}. Since $K_{eq} < K_{neq}$, the periodic potential is always more irrelevant for the nonequilibrium problem, however equations (8), (9) show that the notion of irrelevance should not be taken literally out of equilibrium as the potential can cause inelastic scattering that generates a dissipation rate η, and at long-times a finite temperature T_{eff} [8]. In this paper we will be interested in the pre-thermalized regime where the time is long as compared to microscopic time-scales, while short as compared to the dissipation-rate $1 \ll T_m \ll 1/\eta$ so that η and T_{eff} may be neglected. Since for weak quenches $\eta \sim g^2(K_0 - K)^4$ [9], this regime can be quite large. Eq. (6) also shows that even in the pre-thermalized regime, there is a crossover from an intermediate time dynamics where the physics is determined by the initial wave-function (and hence the initial Luttinger parameter K_0) and a long time time dynamics determined by K_{neq}. This can result in a situation where perturbation theory in g is violated at intermediate times when $\frac{\gamma^2 K_0}{4} < 2$ (*i.e.* g is a relevant perturbation in the initial state). We show below how this happens, and eventually present a RG treatment for the correlator in the regime where we are at weak-coupling at all times. In addition, we will study the system close to the critical point $K_{neq} = 2 + \delta, \forall \delta \ll 1$ where logarithmic corrections to scaling arise.

We evaluate $R = R^{(0)} + R^{(1)} + R^{(2)}$ perturbatively upto quadratic order in the periodic potential ($R^{(i)}$ being of $\mathcal{O}(g^i)$). $R^{(1)}$ at a time T_m after the quench is [29]

$$R^{(1)}(x_1 T_m, x_2 T_m) = \frac{igu}{\alpha^2} \int_{-\infty}^{\infty} dx' \int_0^{T_m} dt' \sum_{c=\pm} sgn(-c)$$
$$\langle \cos\left(\gamma \phi_c(x't') - \frac{\gamma}{2}\phi(x_1 T_m) - \frac{\gamma}{2}\phi(x_2 T_m)\right)\rangle \qquad (11)$$

The behavior of $R^{(1)}$ for several different quench protocols is shown in Fig. 1. We find that outside the light-cone, $T_m < r/2$, the correlator is non-zero. Moreover, for a quench where the initial state is such that the cosine potential is relevant ($\frac{\gamma^2 K_0}{4} < 2$), R can be parametrically large at these initial times outside the light-cone. When the initial state is one where the potential is irrelevant or marginally irrelevant ($\frac{\gamma^2 K_0}{4} \geq 2$), the correlator is most enhanced on the light-cone. For all these cases, within the light-cone $T_m > r/2$, the correlator reaches a steady-state.

We derive the CS-like differential equation for R by splitting the fields ϕ into slow and fast fields $\phi = \phi_< + \phi_>$ where the fast fields have a large weight at short wavelengths, and therefore oscillate rapidly in time. We integrate out the fast fields, and rescale the cut-off, position and time. Such a procedure within the real-time Keldysh approach has been employed for the sine-Gordon model both for steady-state [8, 9, 33] and transient behavior [15], where in each case the β function was derived. Here we generalize this approach to the study of a two-point correlation function. This approach will reveal the



FIG. 1. The first order correction to the equal time correlation function $R^{(1)}(r = 100, T_m)$ as a function of the time T_m after the quench. Length and time are measured in units of the cut-off Λ. Three different quenches are considered: a pure lattice quench ($K_0 = K = 2$), lattice and interaction quench for ($K_0 = 1, K = \sqrt{3}$) and ($K_0 = 3, K = \sqrt{3}$). The light-cone is at $T_m = r/2 = 50$. We choose $\gamma = 2$ and all quenches are to the critical point $K_{neq} = 2$.

existence of a scaling regime where the correlator at long times and distances is related to the correlator at short times and distances (and a renormalized coupling), where the latter may be evaluated readily within perturbation theory.

To leading order ($g = 0$) we find

$$R^{(0)}(r,T_m) = R^{(0)}_<(r,T_m)\left[1 - \frac{d\Lambda}{\Lambda}\gamma_{an,0}(r,T_m)\right] \quad (12)$$

where $R_<$ is the correlator of the slow fields, while R is the correlator for all the fields. The above implies the following differential equation (defining $\Lambda = \Lambda_0/l$, Λ_0 being the bare cut-off, and T_{m0} is the time after the quench), $\left[\frac{\partial}{\partial \ln l} - \gamma_{an,0}(r,T_m)\right] R^{(0)}\left(\frac{r\Lambda_0}{l}, \frac{T_{m0}\Lambda_0}{l}\right) = 0$ where (in units of Λ)

$$\begin{aligned}\gamma_{an,0}(r,T_m) = \frac{1}{2}\Bigg[&K_{neq}\frac{r^2}{1+r^2} + \frac{K_{tr}}{1+(2T_m)^2}\\ &-\frac{K_{tr}}{2}\left\{\frac{1}{1+(2T_m+r)^2} + \frac{1}{1+(2T_m-r)^2}\right\}\Bigg]\end{aligned} \quad (13)$$

Eq. (13) shows that there are two scaling limits where $\gamma_{an,0}$ becomes independent of r, T_m. One is within the light-cone $T_m \gg r \gg 1$ where $\gamma_{an,0} = K_{neq}/2$, and the second is on the light-cone $r = 2T_m, r \gg 1$ where $\gamma_{an,0} = (K_{neq} - K_{tr}/2)/2$. We now discuss the correction to R to next order in g where logarithmic corrections arise when $K_{neq} = 2 + \delta\ \forall\delta \ll 1$.

At next order, in terms of slow and fast fields, we find [29],

$$\begin{aligned}R^{(1)}(r,T_m) = R^{(1)}_<(r,T_m)\Bigg[&1 + 2\frac{d\Lambda}{\Lambda} - 2K_{neq}\frac{d\Lambda}{\Lambda}\\ &+\gamma_{an,0}(r,T_m)\frac{d\Lambda}{\Lambda}\Bigg] + 2g\pi I_C(r,T_m)\frac{d\Lambda}{\Lambda}R^0_<(r,t)\end{aligned} \quad (14)$$

To quadratic order, the correction $R^{(2)}$ leads to the β function which has been discussed above. Eqns. (12), (14) and the β function imply the following CS like differential equation for R,

$$\begin{aligned}&\left[\frac{\partial}{\partial \ln l} + \beta(g_i)\frac{\partial}{\partial g_i} - \gamma_{an,0} + 2\pi g I_C\right]\\ &\times R\left[\frac{r\Lambda_0}{l}, \frac{\Lambda_0 T_{m0}}{l}, g_i(l)\right] = 0\end{aligned} \quad (15)$$

where $g_i = g, \delta, \eta, T_{eff}$. We now discuss the solution to Eq. (15) for two cases, one is for the pure lattice quench and the second for a simultaneous lattice and interaction quench. We will present results in the pre-thermalized regime $T_{m0} \ll 1/\eta$ where η, T_{eff} may be neglected.

Pure lattice quench $K_0 = K$ and $K_{eq} = 2 + \delta\ \forall\delta \ll 1$: Here we find,

$$\begin{aligned}I_C(r,T_m) = &\frac{r^2+1}{4+r^2}\\ &-\left(\frac{1+r^2}{1+T_m^2}\right)\left[\frac{r^2 - 4T_m^2 + 4 - 8T_m^2}{\left\{r^2 - 4T_m^2 + 4\right\}^2 + 64T_m^2}\right]\end{aligned} \quad (16)$$

Note that $I_C(T_m = 0) = 0$ as the lattice has not had time to affect the correlator. Eq. (16) shows the appearance of scaling in two cases, one is within the light-cone where,

$$I_C(r, T_m \gg 1, T_m \gg r) = 1 + \mathcal{O}\left(\frac{1}{r^2}, \frac{r^2}{T_m^4}\right) \quad (17)$$

and the other is on the light-cone,

$$I_C(r, T_m \gg 1, 2T_m = r) = \frac{3}{2} + \mathcal{O}\left(\frac{1}{r^2}\right) \quad (18)$$

Eq. (13), (16) also show that scaling is valid until (restoring units) $l \sim \Lambda_0 r$. In the scaling limit, I_K in Eq. (7) is $I_K(T_m \gg 1) = \pi\left[1 - \frac{7}{8T_m^2} + \ldots\right]$. Thus the solution of the CS equation (15) in the two scaling limits where $I_C, \gamma_{an,0}, I_K$ are constants in time and position is,

$$\begin{aligned}&R\left(\Lambda_0 r, \Lambda_0 T_{m0}, g_0\right) = e^{-\int_{g_0}^{g(l)} dg' \frac{\gamma_{an}(g')}{\beta(g')}}\\ &\times R\left(\frac{r\Lambda_0}{l}, \frac{T_{m0}\Lambda_0}{l}, g(l)\right)\end{aligned} \quad (19)$$

where $\gamma_{an} = 1 + \frac{\delta}{2} - 2\pi g I_C$ and the β function is $\frac{dg}{d\ln l} = -g\delta, \frac{d\delta}{d\ln l} = -(2\pi g)^2$. Note that the critical point corresponding to the $S = 1/2$ Heisenberg chain corresponds to $\delta = 2\pi g$.

The correlation function within the light-cone is obtained by taking $T_m \gg r$, and integrating upto $l = \Lambda_0 r$,

$$R\left(\Lambda_0 r, \frac{T_{m0}}{r} \gg 1, g_0\right) = e^{-\int_{g_0}^{g(\Lambda_0 r)} dg' \frac{\gamma_{an}(g')}{\beta(g')}}$$
$$\times R\left(\frac{r\Lambda_0}{l} = 1, \frac{T_{m0}}{r} \gg 1, g(l)\right) \tag{20}$$

where for $l \gg 1$, $g(l)$ being negligibly small, the correlator on the r.h.s. may be evaluated within perturbation theory. Moreover, $R\left(\frac{r\Lambda_0}{l} = 1, \frac{T_{m0}}{r} \gg 1\right) = \mathcal{O}(1)$ as this is a short-distance correlator. The solution of Eq. (20) at the critical point $\delta{=}2\pi g$ is found to be

$$R(r, T_m \gg r) \simeq \frac{(\ln r)^{I_C - 1/2}}{r} = \frac{\sqrt{\ln r}}{r} \tag{21}$$

In contrast the correlator on the light-cone is

$$R\left(\Lambda_0 r, 2T_{m0} = r, g_0\right) = e^{-\int_{g_0}^{g(\Lambda_0 r)} dg' \frac{\gamma_{an}(g')}{\beta(g')}}$$
$$\times R\left(\frac{r\Lambda_0}{l} = 1, 2T_{m0} = l, g(l)\right) \tag{22}$$

where the r.h.s is again a short distance correlator and therefore of $\mathcal{O}(1)$. Due to Eq. (18), this leads to

$$R(r = 2T_m) = \frac{\ln r}{r} \tag{23}$$

Eq. (21), (23) are the two main results of this section. Thus in the pre-thermalized regime the result Eq. (21) within the light-cone for a pure lattice quench is the same as in the ground state of the Heisenberg chain [25–27]. In contrast, the quench leads to qualitatively new scaling behavior for spatial separations on the light-cone (Eq. (23)). Note that for a pure lattice quench, no dissipative effects are generated at long times to $\mathcal{O}(g^2)$ [9], extending the regime of validity of the pre-thermalized regime.

Simultaneous lattice and interaction quench ($K_0 \neq K$ and $K_{neq} = 2 + \delta, \forall \delta \ll 1$): This quench corresponds to a final Hamiltonian whose ground state can be in the gapped phase. However due to the quench, since $K_{neq} > K_{eq}$, the periodic potential is more irrelevant (though it can give rise to inelastic scattering) [34]. Here we find that I_K in Eq. (7) is $I_K(T_m, K_{neq} = 2){=}\pi\left[c_1 - c_2 \sin\left(\pi(K_{eq} - K_{neq})\right)\ln\left(\min\left[T_m, 1/\eta\right]\right)\right]$ where $c_{1,2}$ are $\mathcal{O}(1)$ and depend on the initial Luttinger parameter K_0. When $K_0{=}K$, $c_1{=}1$. In what follows we will consider a prethermalized regime where $T_m < 1/\eta$. We will also assume, $|K_{eq} - K_{neq}| \ln T_m \ll 1$. In this case, $I_K \simeq \pi c_1$ is a constant in time. The β function in the vicinity of $K_{neq} = 2 + \delta$ becomes $\frac{dg}{d\ln l} = -g\delta, \frac{d\delta}{d\ln l} = -g^2 B^2$, where $B = \pi\sqrt{c_1}\frac{\gamma^2 K_0}{4}\left(\frac{16}{\gamma^2 K_0} - 1\right)^{3/4}$. In addition, I_C in Eq. (15) becomes a universal function of the initial Luttinger parameter K_0 and is plotted in Fig. 2.



FIG. 2. Plot of I_C for a simultaneous lattice and interaction quench for points within the light-cone and near the critical point $K_{neq} = 2$ where $K_{eq} = \frac{\gamma^2 K_0}{4}\sqrt{\frac{16}{\gamma^2 K_0} - 1}$.

Using Eq. (20) to solve for the correlation function within the light cone, the correlation function is

$$R\left(\Lambda_0 r, r \ll T_{m0} \ll 1/\eta, g_0\right) \simeq \frac{1}{(\Lambda_0 r)^{1+A/2}}$$
$$\frac{1}{\sqrt{1 - (r\Lambda_0)^{-2A}}}\left[\frac{1 - (\Lambda_0 r)^{-A}}{1 + (\Lambda_0 r)^{-A}}\right]^{\frac{2\pi I_C}{B}} \tag{24}$$

where $A = \sqrt{\delta^2 - g^2 B^2}$. Thus for an interaction and lattice quench, and for $A \ln r \gg 1$, the correlators decay as a power-law with exponent $(1 + A/2)$. This exponent is not the same as in a quadratic Luttinger liquid theory $(1 + \delta/2)$. In the vicinity of the nonequilibrium critical point $A \to 0$, logarithmic corrections are obtained for $A \ll A \ln r \ll 1$, where

$$R \sim \frac{1}{r}(\ln r)^{\frac{2\pi I_C}{B} - \frac{1}{2}} \tag{25}$$

These corrections are significantly different from that near the equilibrium critical point [25–27]. In contrast to the regime within the light-cone discussed above, scaling behavior is lost on the light-cone for a simultaneous lattice and interaction quench because $I_C(r = 2T_m) \sim r^{K_{tr}/2}$ where $K_{tr} = \frac{\gamma^2 K_0}{4} - 2$ [29].

In summary we have explicitly constructed a CS-like differential equation (15) for a two-point correlation function for a quench in a spin-chain and found that for a lattice quench, scaling holds in two regimes. One is for spatial separations within the light-cone (Eq. (21)), and the second is for spatial separations on the light-cone (Eq. (23)). For a simultaneous lattice and interaction quench, the scaling on the light-cone is lost, but still holds within the light-cone (Eq. (25)). The correlator is found to reach a quasi-steady-state where it decays in position with new power-laws that cannot be recovered within a purely Luttinger liquid model. These results hold in the pre-thermalized regime where dissipative effects are weak. For $T_m > 1/\eta$ dissipative effects will eventually make these correlators decay exponentially fast in position [8]. How this crossover from power-law to exponential decay occurs in time, and observing it in numerical studies [35] are important open questions.

Acknowledgments: The author is deeply indebted to Fabian Essler for many helpful discussions and for a critical reading of the manuscript. This research was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915 and by NSF-DMR (1004589).

[1] I. Bloch, J. Dalibard, and W. Zwerger, Rev. Mod. Phys. **80**, 885 (2008).

[2] D. Fausti, R. I. Tobey, N. Dean, S. Kaiser, A. Dienst, M. C. Hoffmann, S. Pyon, T. Takayama, H. Takagi, and A. Cavalleri, Science **331**, 189 (2011).

[3] C. L. Smallwood, J. P. Hinton, C. Jozwiak, W. Zhang, J. D. Koralek, H. Eisaki, D.-H. Lee, J. Orenstein, and A. Lanzara, Science **336**, 1137 (2012).

[4] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Rev. Mod. Phys. **83**, 863 (2011).

[5] M. Rigol, Phys. Rev. A **80**, 053607 (2009).

[6] L. F. Santos and M. Rigol, Phys. Rev. E **81**, 036206 (2010).

[7] G. Biroli, C. Kollath, and A. M. Läuchli, Phys. Rev. Lett. **105**, 250401 (2010).

[8] A. Mitra and T. Giamarchi, Phys. Rev. Lett. **107**, 150602 (2011).

[9] A. Mitra and T. Giamarchi, Phys. Rev. B **85**, 075117 (2012).

[10] A. Riera, C. Gogolin, and J. Eisert, Phys. Rev. Lett. **108**, 080402 (2012).

[11] J. Berges, S. Borsányi, and C. Wetterich, Phys. Rev. Lett. **93**, 142002 (2004).

[12] M. Moeckel and S. Kehrein, Phys. Rev. Lett. **100**, 175702 (2008).

[13] M. Eckstein and M. Kollar, Phys. Rev. Lett. **100**, 120404 (2008).

[14] M. Heyl, A. Polkovnikov, and S. Kehrein, arXiv:1206.2505 (unpublished).

[15] A. Mitra, Phys. Rev. Lett. **109**, 260601 (2012).

[16] M. A. Cazalilla, Phys. Rev. Lett. **97**, 156403 (2006).

[17] T. Barthel and U. Schollwöck, Phys. Rev. Lett. **100**, 100601 (2008).

[18] J.-S. Caux and R. M. Konik, Phys. Rev. Lett. **109**, 175301 (2012).

[19] F. H. L. Essler, S. Evangelisti, and M. Fagotti, Phys. Rev. Lett. **109**, 247206 (2012).

[20] P. Calabrese, F. H. L. Essler, and M. Fagotti, Phys. Rev. Lett. **106**, 227203 (2011).

[21] D. Iyer and N. Andrei, Phys. Rev. Lett. **109**, 115304 (2012).

[22] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, Phys. Rev. Lett. **98**, 050405 (2007).

[23] M. Kolodrubetz, B. K. Clark, and D. A. Huse, Phys. Rev. Lett. **109**, 015701 (2012).

[24] E. G. D. Torre, E. Demler, and A. Polkovnikov, arXiv:1211.5145 (unpublished).

[25] R. R. P. Singh, M. E. Fisher, and R. Shankar, Phys. Rev. B **39**, 2562 (1989).

[26] I. Affleck, J. Phys. A: Math. Gen. **31**, 4573 (1998).

[27] V. Barzykin and I. Affleck, J. Phys. A: Math. Gen. **32**, 867 (1999).

[28] T. Giamarchi, *Quantum Physics in One Dimension*, Oxford University Press, Oxford (2004).

[29] See Supplementary material for details of the quadratic theory, evaluation of $R^{(1)}$, derivation of the β function, derivation of the Callan-Symanzik like differential equation, and its solution.

[30] A. Kamenev, *Field Theory of Non-Equilibrium Systems*, Cambridge University Press, Cambridge (2011).

[31] A. Iucci and M. A. Cazalilla, Phys. Rev. A **80**, 063619 (2009).

[32] P. Calabrese and J. Cardy, Phys. Rev. Lett. **96**, 136801 (2006).

[33] E. G. Dalla Torre, E. Demler, T. Giamarchi, and E. Altman, Phys. Rev. B **85**, 184302 (2012).

[34] The RG equations show that after a quench V_{sg} may be a weak coupling even though it is a strong coupling for the ground state of the final Hamiltonian.

[35] C. Karrasch, J. Rentrop, D. Schuricht, and V. Meden, Phys. Rev. Lett. **109**, 126406 (2012).

SUPPLEMENTARY MATERIAL: CORRELATION FUNCTIONS AFTER A QUANTUM QUENCH IN A SPIN-CHAIN IN THE PRE-THERMALIZED REGIME

Model

The model we will consider is that of an initial quench from $K_0 \to K$, with the cosine-potential $V = -\frac{gu}{\alpha^2}\int dx \cos(\gamma\phi)$ also switched on suddenly (at the same time as the quench). We use a bosonization prescription where,

$$\phi(x) = -(N_R + N_L)\frac{\pi x}{L} - \frac{i\pi}{L}\sum_{p\neq 0}\left(\frac{L|p|}{2\pi}\right)^{1/2}\frac{1}{p}e^{-\alpha|p|/2-ipx}\left(b_p^\dagger + b_{-p}\right), \tag{26}$$

$$\theta(x) = (N_R - N_L)\frac{\pi x}{L} + \frac{i\pi}{L}\sum_{p\neq 0}\left(\frac{L|p|}{2\pi}\right)^{1/2}\frac{1}{|p|}e^{-\alpha|p|/2-ipx}\left(b_p^\dagger - b_{-p}\right). \tag{27}$$

and $\Lambda = \alpha^{-1}$ is an ultra-violet cutoff.

Thus the initial Hamiltonian for $t \leq 0$ is,

$$\begin{aligned} H_i &= \frac{u_0}{2\pi}\int dx\left[K_0\left(\pi\Pi(x)\right)^2 + \frac{1}{K_0}\left(\partial_x\phi(x)\right)^2\right] \\ &= \sum_{p\neq 0} u_0|p|\eta_p^\dagger\eta_p \end{aligned} \tag{28}$$

while the time evolution from $t > 0$ is due to

$$H_f = H_{f0} + V \tag{29}$$

$$\begin{aligned} H_{f0} &= \frac{u}{2\pi}\int dx\left[K\left(\pi\Pi(x)\right)^2 + \frac{1}{K}\left(\partial_x\phi(x)\right)^2\right] \\ &= \sum_{p\neq 0} u|p|\gamma_p^\dagger\gamma_p \end{aligned} \tag{30}$$

$$V = -\frac{gu}{\alpha^2}\int dx\cos(\gamma\phi) \tag{31}$$

Note that for $L \to \infty$, the zero modes in Eqns. 26, 27 may be dropped. For finite-L, if one assumes that the quench connects the same zero-mode sectors of the initial and final Hamiltonian, they will again not play a role in the dynamics.

To preserve Galilean invariance (which is not necessary for the formalism), we assume $u = v_F/K, u_0 = v_F/K_0$. Note that,

$$\begin{pmatrix} b_p \\ b_{-p}^\dagger \end{pmatrix} = \begin{pmatrix} \cosh\beta & -\sinh\beta \\ -\sinh\beta & \cosh\beta \end{pmatrix}\begin{pmatrix} \gamma_p \\ \gamma_{-p}^\dagger \end{pmatrix} \tag{32}$$

$$\begin{pmatrix} b_p \\ b_{-p}^\dagger \end{pmatrix} = \begin{pmatrix} \cosh\beta_0 & -\sinh\beta_0 \\ -\sinh\beta_0 & \cosh\beta_0 \end{pmatrix}\begin{pmatrix} \eta_p \\ \eta_{-p}^\dagger \end{pmatrix} \tag{33}$$

where $e^{-2\beta_0} = K_0, e^{-2\beta} = K$.

Let us define the functions

$$f(pt) = \cos(u\mid p\mid t)\cosh\beta_0 - i\sin(u\mid p\mid t)\cosh(2\beta - \beta_0) \tag{34}$$

$$g(pt) = \cos(u\mid p\mid t)\sinh\beta_0 + i\sin(u\mid p\mid t)\sinh(2\beta - \beta_0) \tag{35}$$

which determine the time-evolution after the quench ($t > 0$) for $g = 0$,

$$b_p^\dagger(t) + b_{-p}(t) = \left(f^*(pt) - g(pt)\right)\eta_p^\dagger(0) + \left(f(pt) - g^*(pt)\right)\eta_{-p}(0) \tag{36}$$

$$b_p^\dagger(t) - b_{-p}(t) = \left(f^*(pt) + g(pt)\right)\eta_p^\dagger(0) - \left(f(pt) + g^*(pt)\right)\eta_{-p}(0) \tag{37}$$

The Keldysh action is,

$$Z_K = Tr\left[\rho(t)\right] = Tr\left[e^{-iH_ft}|\psi_i\rangle\langle\psi_i|e^{iH_ft}\right] \tag{38}$$

$$= \int \mathcal{D}\left[\phi_{cl}, \phi_q\right]e^{i(S_0 + S_{sg})} \tag{39}$$

where S_0 is the quadratic part which describes the physics in the absence of a periodic potential. In particular at a time t after the quench (note that the fields ϕ are real),

$$S_0 = \frac{1}{2}\int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 \int_0^t dt_1 \int_0^t dt_2 \left(\phi_{cl}(1) \;\; \phi_q(1)\right) \begin{pmatrix} 0 & G_A^{-1}(1,2) \\ G_R^{-1}(1,2) & -\left[G_R^{-1} G_K G_A^{-1}\right](1,2) \end{pmatrix} \begin{pmatrix} \phi_{cl}(2) \\ \phi_q(2) \end{pmatrix} \tag{40}$$

where $1 = (x_1, t_2), 2 = (x_2, t_2)$ and

$$\phi_{cl,q} = \frac{\phi_- \pm \phi_+}{\sqrt{2}} \tag{41}$$

where

$$\left[G^{R,A}(1,2)\right]^{-1} = -\delta(x_1 - x_2)\delta(t_1 - t_2)\frac{1}{\pi K u}\left[\partial^2_{t_1 \pm i\delta} - u^2 \partial^2_{x_1}\right] \tag{42}$$

and,

$$-i\langle \begin{pmatrix} \phi_{cl}(1) \\ \phi_q(1) \end{pmatrix} \left(\phi_{cl}(2) \;\; \phi_q(2)\right)\rangle = \begin{pmatrix} G_K(1,2) & G_R(1,2) \\ G_A(1,2) & 0 \end{pmatrix} \tag{43}$$

Whereas,

$$S_{sg} = \frac{gu}{\alpha^2}\int_{-\infty}^{\infty} dx \int_0^t dt_1 \left[\cos(\gamma\phi_-(1)) \; - \cos(\gamma\phi_+(1))\right] \tag{44}$$

For the quadratic theory after the quench,

$$\begin{aligned} -i\langle\phi(xt_1)\phi(yt_2)\rangle = & -\frac{i}{2}\int_0^{\infty} \frac{dp}{p} e^{-\alpha p}\cos(p(x-y))\left[K_0 \cos(u|p|t_1)\cos(u|p|t_2) + \frac{K^2}{K_0}\sin(u|p|t_1)\sin(u|p|t_2)\right] \\ & -\frac{K}{4}\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right] \end{aligned} \tag{45}$$

The above implies,

$$G^R(xt_1, yt_2) = -i\theta(t_1 - t_2)\langle[\phi(xt_1), \phi(yt_2)]\rangle \tag{46}$$

$$= -\frac{K}{2}\theta(t_1 - t_2)\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right] \tag{47}$$

$$G^A(x-y, t_1-t_2) = G^R(x-y, -(t_1-t_2)) \tag{48}$$

$$= \frac{K}{2}\theta(t_2 - t_1)\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right] \tag{49}$$

and,

$$G^K(xt_1, yt_2) = -i\langle\{\phi(xt_1), \phi(yt_2)\}\rangle \tag{50}$$

$$\begin{aligned} = & -i\frac{K_0}{2}\left(1 + \frac{K^2}{K_0^2}\right)\int_0^{\infty}\frac{dp}{p}e^{-\alpha p}\cos(up(x-y))\cos(u|p|(t_1-t_2)) \\ & -i\frac{K_0}{2}\left(1 - \frac{K^2}{K_0^2}\right)\int_0^{\infty}\frac{dp}{p}e^{-\alpha p}\cos(up(x-y))\cos(u|p|(t_1+t_2)) \end{aligned} \tag{51}$$

Let us define

$$C_{ab}(xt_1, yt_2) = \langle e^{i\gamma\phi_a(xt_1)} e^{-i\gamma\phi_b(yt_2)}\rangle \tag{52}$$

where $a, b = \pm$. Therefore for

$$C_{--}(1,2) = e^{-\frac{\gamma^2}{2}\left[\frac{iG^K(1,1)}{2} + \frac{iG^K(2,2)}{2} - iG^K(1,2)\right]} e^{-iK_{eq} sgn(t_1-t_2)\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right]} \tag{53}$$

$$C_{++}(1,2) = e^{-\frac{\gamma^2}{2}\left[\frac{iG^K(1,1)}{2} + \frac{iG^K(2,2)}{2} - iG^K(1,2)\right]} e^{iK_{eq} sgn(t_1-t_2)\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right]} \tag{54}$$

$$C_{+-}(1,2) = e^{-\frac{\gamma^2}{2}\left[\frac{iG^K(1,1)}{2} + \frac{iG^K(2,2)}{2} - iG^K(1,2)\right]} e^{-iK_{eq}\left[\tan^{-1}\left(\frac{u(t_1-t_2)+x-y}{\alpha}\right) + \tan^{-1}\left(\frac{u(t_1-t_2)-(x-y)}{\alpha}\right)\right]} \tag{55}$$

In general,

$$C_{ab,m}(x_1t_1,x_2t_2) = \langle e^{im\gamma\phi_a(x_1t_1)}e^{-im\gamma\phi_b(x_2t_2)}\rangle \\ e^{-\frac{\gamma^2 m^2}{2}\left[\frac{iG_K(11)}{2}+\frac{iG_K(22)}{2}-iG_K(12)+iaG_A(1,2)+ibG_R(12)\right]} \tag{56}$$

It is convenient to define the following exponents that play an important role in the dynamics,

$$K_{eq} = \frac{\gamma^2 K}{4} \tag{57}$$

$$K_{neq} = \frac{\gamma^2}{8}K_0\left(1+\frac{K^2}{K_0^2}\right) \tag{58}$$

$$K_{tr} = \frac{\gamma^2}{8}K_0\left(1-\frac{K^2}{K_0^2}\right) \tag{59}$$

In terms of these exponents,

$$\begin{aligned}
&-\frac{\gamma^2}{2}\left[\frac{iG^K(1,1)}{2}+\frac{iG^K(2,2)}{2}-iG^K(1,2)\right] \\
&= -K_{neq}\left[\ln\frac{\sqrt{\alpha^2+\{u(t_1-t_2)+x-y\}^2}}{\alpha}+\ln\frac{\sqrt{\alpha^2+\{u(t_1-t_2)-(x-y)\}^2}}{\alpha}\right] \\
&-K_{tr}\left[\ln\sqrt{\frac{\alpha^2+\{u(t_1+t_2)+x-y\}^2}{\alpha^2+(2ut_1)^2}}+\ln\sqrt{\frac{\alpha^2+\{u(t_1+t_2)-(x-y)\}^2}{\alpha^2+(2ut_2)^2}}\right]
\end{aligned} \tag{60}$$

The above shows that the equal-time correlator for positions and times large as compared to the cut-off and for $|r\pm ut|/\alpha \gg 1$ is (setting $\alpha=1$),

$$C_1(rt,0t) = \left[\frac{1}{r^2}\right]^{\frac{\gamma^2 K_0}{4}}\left[|\frac{r^2-(2ut)^2}{r^2(2ut)^2}|\right]^{-K_{tr}} \tag{61}$$

which agrees with Ref. 31 who also point out that for $2ut \ll r$, the correlator decays in position in the same way as in the initial state, but with a time-dependent prefactor (which is $t^{2K_{tr}}$). Exactly at the light-cone $C_1(r=2ut) \simeq \left[\frac{1}{r^2}\right]^{\frac{\gamma^2 K_0}{4}}\left[\frac{1}{r^3}\right]^{-K_{tr}}$. Note that in the main text, the behavior of the correlator $R_{ab} = \langle e^{i\gamma\phi_a(1)/2}e^{-i\gamma\phi_b(2)/2}\rangle$, which follows in a straightforward way from that of C_{ab}, has been discussed.

Derivation of the β function from the action

Let us denote a UV cutoff as

$$\Lambda = \frac{u}{\alpha} \tag{62}$$

We split the fields into slow ($\phi^<$) and fast ($\phi^>$) components [8, 9, 15, 33]

$$\phi_\pm = \phi_\pm^< + \phi_\pm^> \tag{63}$$

and integrate out the fast fields perturbatively. In doing so to $\mathcal{O}(g^2)$, we obtain

$$S = S_0^< + \delta S^< \tag{64}$$

where

$$\begin{aligned}
S_0^< = \int_{-\infty}^{\infty} dR \int_0^{ut/\sqrt{2}} d(uT_m)\frac{1}{2\pi K}&\left[\phi_q^<\left(\partial_R^2-\partial_{uT_m}^2\right)\phi_{cl}^< + \phi_{cl}^<\left(\partial_R^2-\partial_{uT_m}^2\right)\phi_q^< + \frac{\delta u}{u}\phi_q^<\left(\partial_R^2+\partial_{uT_m}^2\right)\phi_{cl}^<\right. \\
&\left. +\frac{\delta u}{u}\phi_{cl}^<\left(\partial_R^2+\partial_{uT_m}^2\right)\phi_q^< - 2\frac{\eta}{u}\phi_q^<\partial_{uT_m}\phi_{cl}^< + i\frac{4\eta T_{eff}}{u^2}\left(\phi_q^<\right)^2\right]
\end{aligned} \tag{65}$$

where $T_{eff}=\eta=0$ initially.

The correction from integrating the fast fields is,

$$\delta S^{<} = \frac{g\Lambda^2}{u}\int_{-\infty}^{\infty} dx \int_0^t dt_1 \left[\cos\gamma\phi_{-}^{<}(1) - \cos\gamma\phi_{+}^{<}(1)\right] e^{-\frac{\gamma^2}{4}\langle\left(\phi_{cl}^{>}(1)\right)^2\rangle} \tag{66}$$

$$+\frac{ig^2\Lambda^4}{2u^2}\int_{-\infty}^{\infty} dx_1 \int_0^t dt_1 \int_{-\infty}^{\infty} dx_2 \int_0^t dt_2 \theta(t_1 - t_2) : \cos\left(\gamma\phi_{-}^{<}(1) - \gamma\phi_{-}^{<}(2)\right) : e^{-\frac{\gamma^2}{2}\langle(\phi_{-}(1)-\phi_{-}(2))^2\rangle}$$

$$\times\left[1 - e^{-\gamma^2\langle\phi_{-}^{>}(1)\phi_{-}^{>}(2)\rangle}\right] \tag{67}$$

$$+\frac{ig^2\Lambda^4}{2u^2}\int_{-\infty}^{\infty} dx_1 \int_0^t dt_1 \int_{-\infty}^{\infty} dx_2 \int_0^t dt_2 \theta(t_2 - t_1) : \cos\left(\gamma\phi_{+}^{<}(1) - \gamma\phi_{+}^{<}(2)\right) : e^{-\frac{\gamma^2}{2}\langle(\phi_{+}(1)-\phi_{+}(2))^2\rangle}$$

$$\times\left[1 - e^{-\gamma^2\langle\phi_{+}^{>}(1)\phi_{+}^{>}(2)\rangle}\right] \tag{68}$$

$$-\frac{ig^2\Lambda^4}{2u^2}\int_{-\infty}^{\infty} dx_1 \int_0^t dt_1 \int_{-\infty}^{\infty} dx_2 \int_0^t dt_2 \{\theta(t_1 - t_2) + \theta(t_2 - t_1)\} : \cos\left(\gamma\phi_{+}^{<}(1) - \gamma\phi_{-}^{<}(2)\right) : e^{-\frac{\gamma^2}{2}\langle(\phi_{+}(1)-\phi_{-}(2))^2\rangle}$$

$$\times\left[1 - e^{-\gamma^2\langle\phi_{+}^{>}(1)\phi_{-}^{>}(2)\rangle}\right] \tag{69}$$

where the fast correlators are related in a simple way to derivatives of the full correlators,

$$G^{>} = d\Lambda\frac{dG}{d\Lambda} \tag{70}$$

Explicit expressions for the fast correlators at equal time are

$$\langle\left(\phi_{cl}^{>}(t)\right)^2\rangle = \frac{d\Lambda}{\Lambda}\left[\frac{K_0}{2}\left(1 + \frac{K^2}{K_0^2}\right) + \frac{K_0}{2}\left(1 - \frac{K^2}{K_0^2}\right)\left\{\frac{1}{1 + \left(2t\Lambda\right)^2}\right\}\right] \tag{71}$$

$$\xrightarrow{t\ll 1/\Lambda} \frac{d\Lambda}{\Lambda}K_0 \tag{72}$$

$$\xrightarrow{t\gg 1/\Lambda} \frac{d\Lambda}{\Lambda}\frac{K_0}{2}\left(1 + \frac{K^2}{K_0^2}\right) \tag{73}$$

For the non-local fast correlators, we have

$$\langle\phi_{cl}^{>}(1)\phi_{cl}^{>}(2)\rangle = \frac{d\Lambda}{\Lambda}\left[\frac{K_0}{4}\left(1 + \frac{K^2}{K_0^2}\right)\left\{\frac{1}{1 + \Lambda^2((t_1 - t_2) + (x_1 - x_2)/u)^2} + \frac{1}{1 + \Lambda^2((t_1 - t_2) - (x_1 - x_2)/u)^2}\right\}\right.$$

$$\left.+\frac{K_0}{4}\left(1 - \frac{K^2}{K_0^2}\right)\left\{\frac{1}{1 + \Lambda^2((t_1 + t_2) + (x_1 - x_2)/u)^2} + \frac{1}{1 + \Lambda^2((t_1 + t_2) - (x_1 - x_2)/u)^2}\right\}\right] \tag{74}$$

$$\langle\phi_{cl}^{>}(1)\phi_{q}^{>}(2)\rangle = -i\frac{K}{2}\frac{d\Lambda}{\Lambda}\theta(t_1 - t_2)\left[\frac{\Lambda((t_1 - t_2) + (x_1 - x_2)/u)}{1 + \Lambda^2((t_1 - t_2) + (x_1 - x_2)/u)^2} + \frac{\Lambda((t_1 - t_2) - (x_1 - x_2)/u)}{1 + \Lambda^2((t_1 - t_2) - (x_1 - x_2)/u)^2}\right] \tag{75}$$

$$\langle\phi_{q}^{>}(1)\phi_{cl}^{>}(2)\rangle = i\frac{K}{2}\frac{d\Lambda}{\Lambda}\theta(t_2 - t_1)\left[\frac{\Lambda((t_1 - t_2) + (x_1 - x_2)/u)}{1 + \Lambda^2((t_1 - t_2) + (x_1 - x_2)/u)^2} + \frac{\Lambda((t_1 - t_2) - (x_1 - x_2)/u)}{1 + \Lambda^2((t_1 - t_2) - (x_1 - x_2)/u)^2}\right] \tag{76}$$

In the next step we define new variables

$$R = \frac{x_1 + x_2}{2} \ \ , T_m = \frac{t_1 + t_2}{2} \tag{77}$$

$$r = x_1 - x_2 \ \ , \tau = t_1 - t_2 \tag{78}$$

Thus

$$\int_{-\infty}^{\infty} dx_1 \int_{-\infty}^{\infty} dx_2 = \int_{-\infty}^{\infty} dR \int_{-\infty}^{\infty} dr \tag{79}$$

$$\int_0^t dt_1 \int_0^t dt_2 = \int_0^{t/\sqrt{2}} dT_m \int_{-2T_m}^{2T_m} d\tau \tag{80}$$

and perform a gradient expansion in R, T_m to obtain,

$$\delta S^{<} = \delta S_g^{<} + \delta S_0^{<} + \delta S_{T_{eff}}^{<} + \delta S_\eta^{<} \tag{81}$$

where

$$\delta S_g^< = \frac{gu}{\alpha^2}\int_{-\infty}^{\infty} dx \int_0^t dt_1 \left[\cos\gamma\phi_-^<(1) - \cos\gamma\phi_+^<(1)\right] e^{-\frac{\gamma^2}{4}\langle\left(\phi_{cl}^>(1)\right)^2\rangle} \tag{82}$$

and

$$\begin{aligned}\delta S_0^< = {} & \frac{g^2\Lambda^4\gamma^2}{2u^2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} dR \int_{-\infty}^{\infty} dr \int_0^{t/\sqrt{2}} dT_m \int_{-2T_m}^{2T_m} d\tau\theta(\tau)\left[\left(r\partial_R\phi_{cl}^<\right)\left(r\partial_R\phi_q^<\right) + \left(\tau\partial_{T_m}\phi_{cl}^<\right)\left(\tau\partial_{T_m}\phi_q^<\right)\right] \\ & \times Im\left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle}F(r,T_m,\tau)\right]\end{aligned} \tag{83}$$

while

$$\begin{aligned}\delta S_{T_{eff}}^< = {} & \frac{ig^2\Lambda^4\gamma^2}{2u^2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} dR \int_{-\infty}^{\infty} dr \int_0^{t/\sqrt{2}} dT_m \int_{-2T_m}^{2T_m} d\tau \left(\phi_q^<(R,T_m)\right)^2 \\ & \times Re\left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle}F(r,T_m,\tau)\right]\end{aligned} \tag{84}$$

and

$$\begin{aligned}\delta S_\eta^< = {} & \frac{g^2\Lambda^4\gamma^2}{2u^2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} dR \int_{-\infty}^{\infty} dr \int_0^{t/\sqrt{2}} dT_m \int_{-2T_m}^{2T_m} d\tau\phi_q^<(R,T_m)\tau\partial_{T_m}\phi_{cl}^<(R,T_m) \\ & \times Im\left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle}F(r,T_m,\tau)\right]\end{aligned} \tag{85}$$

where

$$\begin{aligned}F(r,T_m,\tau) = {} & K_{neq}\left[\frac{1}{1+\Lambda^2\left(\tau+r/u\right)^2} + \frac{1}{1+\Lambda^2\left(\tau-r/u\right)^2}\right] \\ & + K_{tr}\left[\frac{1}{1+\Lambda^2\left(2T_m+r/u\right)^2} + \frac{1}{1+\Lambda^2\left(2T_m-r/u\right)^2}\right] \\ & - iK_{eq}\left[\frac{\Lambda(\tau+r/u)}{1+\Lambda^2\left(\tau+r/u\right)^2} + \frac{\Lambda(\tau-r/u)}{1+\Lambda^2\left(\tau-r/u\right)^2}\right]\end{aligned} \tag{86}$$

whereas

$$\begin{aligned} & e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle} = \left[\frac{1}{\sqrt{1+\Lambda^2(\tau+r/u)^2}}\frac{1}{\sqrt{1+\Lambda^2(\tau-r/u)^2}}\right]^{K_{neq}} \\ & \times\left[\frac{\sqrt{1+\Lambda^2\{2(T_m+\tau/2)\}^2}}{\sqrt{1+\Lambda^2(2T_m+r/u)^2}}\frac{\sqrt{1+\Lambda^2\{2(T_m-\tau/2)\}^2}}{\sqrt{1+\Lambda^2(2T_m-r/u)^2}}\right]^{K_{tr}} \\ & \times e^{-iK_{eq}\left[\tan^{-1}(\Lambda(\tau+r/u))+\tan^{-1}(\Lambda(\tau-r/u))\right]}\end{aligned} \tag{87}$$

where $Re[A] = (A+A^*)/2, Im[A] = (A-A^*)/(2i)$.

Collecting all terms we find,

$$\delta S_g^< = g\Lambda^2\int_{-\infty}^{\infty} d(R/u)\int_0^t dT_m\left[\cos\gamma\phi_-^<(R,T_m) - \cos\gamma\phi_+^<(R,T_m)\right]e^{-\frac{\gamma^2}{4}\langle\left(\phi_{cl}^>(T_m)\right)^2\rangle} \tag{88}$$

$$\delta S_0^< = \frac{g^2\gamma^2}{2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} d(R/u)\int_0^{t/\sqrt{2}} dT_m\left[-I_R(T_m)\left(\partial_{R/u}\phi_{cl}^<\right)\left(\partial_{R/u}\phi_q^<\right) - I_{T_m}(T_m)\left(\partial_{T_m}\phi_{cl}^<\right)\left(\partial_{T_m}\phi_q^<\right)\right] \tag{89}$$

$$\delta S_{T_{eff}}^< = \frac{ig^2\gamma^2\Lambda^2}{2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} d(R/u)\int_0^{t/\sqrt{2}} dT_m\left(\phi_q^<\right)^2 I_{T_{eff}}(T_m) \tag{90}$$

$$\delta S_\eta^< = -\frac{g^2\gamma^2\Lambda}{2}\frac{d\Lambda}{\Lambda}\int_{-\infty}^{\infty} d(R/u)\int_0^{t/\sqrt{2}} dT_m\phi_q^<\left[\partial_{T_m}\phi_{cl}^<\right]I_\eta(T_m) \tag{91}$$

where

$$I_R(T_m) = -\Lambda^4 \int_{-\infty}^{\infty} d(r/u) \int_0^{2T_m} d\tau (r/u)^2 Im \left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle} F(r,T_m,\tau)\right] \tag{92}$$

$$I_{T_m}(T_m) = -\Lambda^4 \int_{-\infty}^{\infty} d(r/u) \int_0^{2T_m} d\tau \tau^2 Im \left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle} F(r,T_m,\tau)\right] \tag{93}$$

$$I_{T_{eff}}(T_m) = \Lambda^2 \int_{-\infty}^{\infty} d(r/u) \int_{-2T_m}^{2T_m} d\tau Re \left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle} F(r,T_m,\tau)\right] \tag{94}$$

$$I_\eta(T_m) = -\Lambda^3 \int_{-\infty}^{\infty} d(r/u) \int_{-2T_m}^{2T_m} d\tau \tau Im \left[e^{-\frac{\gamma^2}{2}\langle[\phi_+(R+r/2,T_m+\tau/2)-\phi_-(R-r/2,T_m-\tau/2)]^2\rangle} F(r,T_m,\tau)\right] \tag{95}$$

$$I_K = I_R - I_{T_m} \tag{96}$$

$$I_u = I_R + I_{T_m} \tag{97}$$

At the next step we rescale the cut-off back to the original value of Λ, and in the process rescale position and time to $R, T_m \to \frac{\Lambda}{\Lambda'}(R, T_m)$. This rescaling is not necessary in expressions for $\delta S^{<}_{0,T_{eff},\eta}$ as they are already of $\mathcal{O}\left(\frac{d\Lambda}{\Lambda}\right)$. We also express everything in dimensionless units of $\bar{R} = \Lambda R/u, \bar{T} = T\Lambda$. Thus to summarize one obtains,

$$\begin{aligned} S_0^{<} = \int_{-\infty}^{\infty} d\bar{R} \int_0^{\frac{t}{\sqrt{2}}\Lambda\left(\frac{\Lambda'}{\Lambda}\right)} d\bar{T}_m \frac{1}{2\pi K} \Big[&\phi_q^{<} \left(\partial_{\bar{R}}^2 - \partial_{\bar{T}_m}^2\right) \phi_{cl}^{<} + \phi_{cl}^{<} \left(\partial_{\bar{R}}^2 - \partial_{\bar{T}_m}^2\right) \phi_q^{<} + \frac{\delta u}{u} \phi_q^{<} \left(\partial_{\bar{R}}^2 + \partial_{\bar{T}_m}^2\right) \phi_{cl}^{<} \\ &+ \frac{\delta u}{u} \phi_{cl}^{<} \left(\partial_{\bar{R}}^2 + \partial_{\bar{T}_m}^2\right) \phi_q^{<} - 2\eta \left(\frac{\Lambda}{\Lambda'}\right) \phi_q^{<} \partial_{\bar{T}_m} \phi_{cl}^{<} + i4\eta T_{eff} \left(\frac{\Lambda}{\Lambda'}\right)^2 \left(\phi_q^{<}\right)^2 \Big] \end{aligned} \tag{98}$$

and

$$\delta S_g^{<} = g \left(\frac{\Lambda}{\Lambda'}\right)^2 \int_{-\infty}^{\infty} d\bar{R} \int_0^{t\Lambda\left(\frac{\Lambda'}{\Lambda}\right)} d\bar{T}_m \left[\cos\gamma\phi_-^{<}(R,T_m) - \cos\gamma\phi_+^{<}(R,T_m)\right] e^{-\frac{\gamma^2}{4}\langle\left(\phi_{cl}^{>}(T_m)\right)^2\rangle} \tag{99}$$

$$\delta S_0^{<} = \frac{g^2\gamma^2}{2} \frac{d\Lambda}{\Lambda} \int_{-\infty}^{\infty} d\bar{R} \int_0^{t\Lambda/\sqrt{2}} d\bar{T}_m \left[-I_R(T_m) \left(\partial_{\bar{R}} \phi_{cl}^{<}\right) \left(\partial_{\bar{R}} \phi_q^{<}\right) - I_{T_m}(T_m) \left(\partial_{\bar{T}_m} \phi_{cl}^{<}\right) \left(\partial_{\bar{T}_m} \phi_q^{<}\right)\right] \tag{100}$$

$$\delta S_{T_{eff}}^{<} = \frac{ig^2\gamma^2}{2} \frac{d\Lambda}{\Lambda} \int_{-\infty}^{\infty} d\bar{R} \int_0^{t\Lambda/\sqrt{2}} d\bar{T}_m \left(\phi_q^{<}\right)^2 I_{T_{eff}}(T_m) \tag{101}$$

$$\delta S_\eta^{<} = -\frac{g^2\gamma^2}{2} \frac{d\Lambda}{\Lambda} \int_{-\infty}^{\infty} d\bar{R} \int_0^{t\Lambda/\sqrt{2}} d\bar{T}_m \phi_q^{<} \left[\partial_{\bar{T}_m} \phi_{cl}^{<}\right] I_\eta(T_m) \tag{102}$$

Since,

$$\frac{\Lambda}{\Lambda'} = e^{d\ln(l)} \tag{103}$$

$$|\frac{d\Lambda}{\Lambda}| = \frac{\Lambda - \Lambda'}{\Lambda} = d\ln(l) \tag{104}$$

Therefore the RG equations in terms of dimensionless variables such as $T_m \to T_m\Lambda, \eta \to \eta/\Lambda, T_{eff} \to T_{eff}/\Lambda$ are

$$\frac{dg}{d\ln l} = g\left[2 - \left(K_{neq} + \frac{K_{tr}}{1 + 4T_m^2}\right)\right] \tag{105}$$

$$\frac{dK^{-1}}{d\ln l} = \frac{\pi g^2\gamma^2}{4} I_K(T_m) \tag{106}$$

$$\frac{1}{Ku}\frac{du}{d\ln l} = \frac{\pi g^2\gamma^2}{4} I_u(T_m) \tag{107}$$

$$\frac{d\eta}{d\ln l} = \eta + \frac{\pi g^2\gamma^2 K}{2} I_\eta(T_m) \tag{108}$$

$$\frac{d(\eta T_{eff})}{d\ln l} = 2\eta T_{eff} + \frac{\pi g^2\gamma^2 K}{4} I_{T_{eff}}(T_m) \tag{109}$$

$$\frac{dT_m}{d\ln l} = -T_m \tag{110}$$

Note that the renormalization of the velocity in Eq. 107 is a minor effect which will be neglected in the main text. To summarize:

$$I_R(T_m\Lambda) = -\int_{-\infty}^{\infty} d\bar{r}\int_0^{2T_m\Lambda} d\bar{\tau}\bar{r}^2 \mathrm{Im}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{111}$$

$$I_{ti}(T_m\Lambda) = -\int_{-\infty}^{\infty} d\bar{r}\int_0^{2T_m\Lambda} d\bar{\tau}\bar{\tau}^2 \mathrm{Im}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{112}$$

$$I_{T_{eff}}(T_m\Lambda) = \int_{-\infty}^{\infty} d\bar{r}\int_{-2T_m\Lambda}^{2T_m\Lambda} d\bar{\tau} \mathrm{Re}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{113}$$

$$I_{\eta}(T_m\Lambda) = -\int_{-\infty}^{\infty} d\bar{r}\int_{-2T_m\Lambda}^{2T_m\Lambda} d\bar{\tau}\bar{\tau} \mathrm{Im}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{114}$$

$$I_u(T_m\Lambda) = -\int_{-\infty}^{\infty} d\bar{r}\int_0^{2T_m\Lambda} d\bar{\tau}\left(\bar{r}^2+\bar{\tau}^2\right) \mathrm{Im}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{115}$$

$$I_K(T_m\Lambda) = -\int_{-\infty}^{\infty} d\bar{r}\int_0^{2T_m\Lambda} d\bar{\tau}\left(\bar{r}^2-\bar{\tau}^2\right) \mathrm{Im}\left[B(\bar{r},T_m\Lambda,\bar{\tau})\right] \tag{116}$$

where $\mathrm{Re}[B] = (B+B^*)/2$, $\mathrm{Im}[B]=(B-B^*)/(2i)$, and

$$B(\bar{r},T_m\Lambda,\bar{\tau}) = C_{+-}(\bar{r},T_m\Lambda,\bar{\tau})F(\bar{r},T_m\Lambda,\bar{\tau}) \tag{117}$$

with $C_{+-}(\bar{r},T_m\Lambda,\bar{\tau}) = \langle e^{i\phi_+(\bar{r},\bar{\tau}+T_m\Lambda/2)}e^{-i\phi_-(0,\bar{\tau}-T_m\Lambda/2)}\rangle$. This quantity within leading order in perturbation theory is,

$$\begin{aligned} &C_{+-}(\bar{r},T_m\Lambda,\bar{\tau}) = \left[\frac{1}{\sqrt{1+(\bar{\tau}+\bar{r})^2}}\frac{1}{\sqrt{1+(\bar{\tau}-\bar{r})^2}}\right]^{K_{neq}} \\ &\left[\frac{\sqrt{1+\{2(T_m\Lambda+\bar{\tau}/2)\}^2}}{\sqrt{1+(2T_m\Lambda+\bar{r})^2}}\frac{\sqrt{1+\{2(T_m\Lambda-\bar{\tau}/2)\}^2}}{\sqrt{1+(2T_m\Lambda-\bar{r})^2}}\right]^{K_{tr}} \\ &\times e^{-iK_{eq}\left[\tan^{-1}(\bar{\tau}+\bar{r})+\tan^{-1}(\bar{\tau}-\bar{r})\right]} \end{aligned} \tag{118}$$

while F is given by

$$\begin{aligned} &F(\bar{r},T_m\Lambda,\bar{\tau}) = K_{neq}\left[\frac{1}{1+(\bar{\tau}+\bar{r})^2}+\frac{1}{1+(\bar{\tau}-\bar{r})^2}\right] \\ &+K_{tr}\left[\frac{1}{1+(2T_m\Lambda+\bar{r})^2}+\frac{1}{1+(2T_m\Lambda-\bar{r})^2}\right] \\ &-iK_{eq}\left[\frac{\bar{\tau}+\bar{r}}{1+(\bar{\tau}+\bar{r})^2}+\frac{\bar{\tau}-\bar{r}}{1+(\bar{\tau}-\bar{r})^2}\right] \end{aligned} \tag{119}$$

Perturbative evaluation of equal time correlation function

Let us study the following two-point function to next to leading order in perturbation theory, where

$$R(x_1t,x_2t) = 4\langle\cos\left(\frac{\gamma\phi(x_1t)}{2}\right)\cos\left(\frac{\gamma\phi(x_2t)}{2}\right)\rangle \tag{120}$$

$$= R^{(0)}+R^{(1)}+\dots \tag{121}$$

where

$$R^{(0)}(x_1t,x_2t) = 2e^{-\frac{\gamma^2}{8}\left[\frac{iG_K(11)}{2}+\frac{iG_K(22)}{2}-iG_K(12)\right]} \tag{122}$$

Using Eq. 60 we get

$$R^{(0)}(x_1t, x_2t) = 2\left[\left(\frac{1}{\sqrt{1+\frac{(x_1-x_2)^2}{\alpha^2}}}\right)^{2K_{neq}}\left(\frac{\sqrt{1+\frac{(2ut+x_1-x_2)^2}{\alpha^2}}\sqrt{1+\frac{(2ut-x_1+x_2)^2}{\alpha^2}}}{1+\frac{(2ut)^2}{\alpha^2}}\right)^{-K_{tr}}\right]^{1/4} \tag{123}$$

The above may be rewritten to give in the scaling limit, Eq. 5 in the main text.

To next order the equal time correlator is,

$$R^{(1)}(x_1t, x_2t) = \frac{igu}{\alpha^2}\int_{-\infty}^{\infty}dx'\int_0^t dt'\sum_{c=\pm} sgn(-c)4\langle\cos\left(\gamma\phi_c(x't')\right)\cos\left(\frac{\gamma}{2}\phi(x_1t)\right)\cos\left(\frac{\gamma}{2}\phi(x_2t)\right)\rangle \tag{124}$$

On evaluating the expectation value, one finds,

$$\begin{aligned}R^{(1)}(x_1t, x_2t) &= \left(\frac{2gu}{\alpha^2}\right)e^{\frac{\gamma^2}{8}\left[\frac{iG_K(11)}{2}+\frac{iG_K(22)}{2}-iG_K(12)\right]}\\ &\int_{-\infty}^{\infty}dx'\int_0^t dt'\sin\left[\frac{\gamma^2K}{8}\left\{\tan^{-1}\left(\frac{u(t-t')+x_1-x'}{\alpha}\right)+\tan^{-1}\left(\frac{u(t-t')-(x_1-x')}{\alpha}\right)\right\}\right.\\ &\left.+\frac{\gamma^2K}{8}\left\{\tan^{-1}\left(\frac{u(t-t')+x_2-x'}{\alpha}\right)+\tan^{-1}\left(\frac{u(t-t')-(x_2-x')}{\alpha}\right)\right\}\right]\\ &e^{-\frac{\gamma^2}{4}\left[\frac{iG_K(1'1')}{2}+\frac{iG_K(11)}{2}-iG_K(1'1)+\frac{iG_K(1'1')}{2}+\frac{iG_K(22)}{2}-iG_K(1'2)\right]}\end{aligned} \tag{125}$$

The above expression for $R^{(1)}$ for different quench protocols is discussed and plotted in Fig. 1 of the main text.

At short times

$$\begin{aligned}R^{(1)}(x_1, x_2; t\Lambda \ll 1) &\rightarrow g\left(\frac{\gamma^2K}{4}\right)(\Lambda t)^2\int_{-\infty}^{\infty}\frac{dx'}{\alpha}\left[\frac{1}{1+(\frac{x_1-x'}{\alpha})^2}+\frac{1}{1+(\frac{x_2-x'}{\alpha})^2}\right]\\ &\times\left(\frac{\sqrt{1+(\frac{x_1-x_2}{\alpha})^2}}{\left[1+(\frac{x'-x_1}{\alpha})^2\right]\left[1+(\frac{x'-x_2}{\alpha})^2\right]}\right)^{\frac{\gamma^2K_0}{8}}\end{aligned} \tag{126}$$

$$R^{(1)}(x_1, x_2; t\Lambda \ll 1) \xrightarrow{|x_1-x_2|\gg 1} g\left(\frac{\gamma^2K}{4}\right)(\Lambda t)^2 R^{(0)}(|x_1-x_2|)c_1\left[\frac{\gamma^2K_0}{8}, |x_1-x_2| \gg 1\right] \tag{127}$$

where $c_1\left[\theta, |x_1-x_2|\right] = \int_{-\infty}^{\infty}dx\frac{|x_1-x_2|^{2\theta}}{[1+x^2]^{1+\theta}[1+(x-(x_1-x_2))^2]^{\theta}}$ is independent of $|x_1-x_2|$ for $|x_1-x_2| \gg 1$

Derivation of a Callan-Symanzik-like differential equation out of equilibrium

In order to derive a CS-like differential equation for the equal-time correlation function

$$\begin{aligned}R(x_1t, x_2t) &= 4\langle\cos\left(\frac{\gamma\phi(x_1t)}{2}\right)\cos\left(\frac{\gamma\phi(x_2t)}{2}\right)\rangle &(128)\\ &= R^{(0)}+R^{(1)}+\ldots &(129)\end{aligned}$$

we split the fields ϕ, into slow and fast fields, and integrate over the fast fields. In doing so, for example, $R^{(0)}$ may be written as,

$$R^{(0)}(x_1t, x_2t) = R^0_<(x_1t, x_2t)e^{-\frac{d\Lambda}{2\Lambda}\left[K_{neq}\frac{(x_1-x_2)^2\Lambda^2/u^2}{1+(x_1-x_2)^2\Lambda^2/u^2}+\frac{K_{tr}}{1+(2t\Lambda)^2}-\frac{K_{tr}}{2}\left\{\frac{1}{1+\Lambda^2(2t+(x_1-x_2)/u)^2}+\frac{1}{1+\Lambda^2(2t-(x_1-x_2)/u)^2}\right\}\right]} \tag{130}$$

Then we rescale the cut-off, and also rescale position and time. In doing so, the above expression does not change. Expanding the above expression to $\mathcal{O}\left(\frac{d\Lambda}{\Lambda}\right)$, and noting that

$$\Lambda\frac{R^{(0)}(\Lambda)-R^{(0)}_<(\Lambda-d\Lambda)}{d\Lambda} = \Lambda\frac{\partial R^{(0)}}{\partial\Lambda} \tag{131}$$

we obtain the CS-like equation to leading order,

$$\begin{aligned}\left[\frac{\partial}{\partial\ln\Lambda}+\gamma_{an,0}(x_1t,x_2t)\right]R^{(0)}(\Lambda)=0\\ \left[\frac{\partial}{\partial\ln l}-\gamma_{an,0}(x_1t,x_2t)\right]R^{(0)}\left(\frac{\Lambda_0}{l}\right)=0\end{aligned} \tag{132}$$

where Λ_0 is the bare cut-off and,

$$\begin{aligned}\gamma_{an,0}(x_1t,x_2t)=&\,\frac{1}{2}\left[K_{neq}\frac{(x_1-x_2)^2\Lambda^2/u^2}{1+(x_1-x_2)^2\Lambda^2/u^2}+\frac{K_{tr}}{1+(2t\Lambda)^2}\right.\\ &\left.-\frac{K_{tr}}{2}\left\{\frac{1}{1+\Lambda^2(2t+(x_1-x_2)/u)^2}+\frac{1}{1+\Lambda^2(2t-(x_1-x_2)/u)^2}\right\}\right]\end{aligned} \tag{133}$$

Some limiting expressions for $\gamma_{an,0}$ are

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\ll 1)=\frac{1}{2}\left(K_{neq}+K_{tr}\right)=\frac{\gamma^2K_0}{8} \tag{134}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t\gg|x_1-x_2|)=\frac{K_{neq}}{2} \tag{135}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t\ll|x_1-x_2|)=\frac{K_{neq}}{2} \tag{136}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t=|x_1-x_2|)=\frac{1}{2}\left(K_{neq}-\frac{K_{tr}}{2}\right) \tag{137}$$

Now for $R^{(1)}$ we find,

$$R^{(1)}(x_1t,x_2t)=4\mathrm{Tr}\left[e^{i(S_0^<+S_0^>)}iS_{sg}(\phi_<+\phi_>)\cos\left(\frac{\gamma\phi^<(x_1t)+\gamma\phi^>(x_1t)}{2}\right)\cos\left(\frac{\gamma\phi^<(x_2t)+\gamma\phi^>(x_2t)}{2}\right)\right] \tag{138}$$

After some algebra we obtain,

$$\begin{aligned}R^{(1)}(x_1t,x_2t)=&\,\frac{igu}{\alpha^2}\int_{-\infty}^{\infty}dx'\int_0^t dt'\sum_{c=\pm}sgn(-c)\\ &\langle\cos\left(\gamma\phi_c^<(x't')-\frac{\gamma}{2}\phi^<(x_1t)-\frac{\gamma}{2}\phi^<(x_2t)\right)\rangle\\ &\times\langle\cos\left(\gamma\phi_c^>(x't')-\frac{\gamma}{2}\phi^>(x_1t)-\frac{\gamma}{2}\phi^>(x_2t)\right)\rangle\end{aligned} \tag{139}$$

$$\begin{aligned}=&\,\frac{igu}{\alpha^2}e^{\frac{d\Lambda}{\Lambda}\gamma_{an,0}(x_1,x_2,t)}\int_{-\infty}^{\infty}dx'\int_0^t dt'\sum_{c=\pm}sgn(-c)e^{-\frac{d\Lambda}{\Lambda}\delta(x_1,x_2,t,x',t')}\\ &\times\langle\cos\left(\gamma\phi_c^<(x't')-\frac{\gamma}{2}\phi^<(x_1t)-\frac{\gamma}{2}\phi^<(x_2t)\right)\rangle\end{aligned} \tag{140}$$

where

$$\begin{aligned}\delta(x_1,x_2,t,x',t')=&\left[2K_{neq}+\frac{K_{tr}}{1+(2t'\Lambda)^2}+\frac{K_{tr}}{1+(2t\Lambda)^2}\right.\\ &-\frac{K_{neq}}{2}\left\{\frac{1}{1+\Lambda^2(t-t'+(x_1-x')/u)^2}+\frac{1}{1+\Lambda^2(t-t'-(x_1-x')/u)^2}\right\}\\ &-\frac{K_{tr}}{2}\left\{\frac{1}{1+\Lambda^2(t+t'+(x_1-x')/u)^2}+\frac{1}{1+\Lambda^2(t+t'-(x_1-x')/u)^2}\right\}\\ &-\frac{K_{neq}}{2}\left\{\frac{1}{1+\Lambda^2(t-t'+(x_2-x')/u)^2}+\frac{1}{1+\Lambda^2(t-t'-(x_2-x')/u)^2}\right\}\\ &-\frac{K_{tr}}{2}\left\{\frac{1}{1+\Lambda^2(t+t'+(x_2-x')/u)^2}+\frac{1}{1+\Lambda^2(t+t'-(x_2-x')/u)^2}\right\}\\ &-ic\frac{K_{eq}}{2}\left\{\frac{\Lambda(t-t'+(x_1-x')/u)}{1+\Lambda^2(t-t'+(x_1-x')/u)^2}+\frac{\Lambda(t-t'-(x_1-x')/u)}{1+\Lambda^2(t-t'-(x_1-x')/u)^2}\right\}\\ &\left.-ic\frac{K_{eq}}{2}\left\{\frac{\Lambda(t-t'+(x_2-x')/u)}{1+\Lambda^2(t-t'+(x_2-x')/u)^2}+\frac{\Lambda(t-t'-(x_2-x')/u)}{1+\Lambda^2(t-t'-(x_2-x')/u)^2}\right\}\right]\end{aligned} \tag{141}$$

Rescaling the cut-off and correspondingly the position and time, we get

$$\begin{aligned} R^{(1)}(x_1-x_2,t) =& R_<^{(1)}(x_1-x_2,t)\left[1+2\frac{d\Lambda}{\Lambda}-2K_{neq}\frac{d\Lambda}{\Lambda}+\gamma_{an,0}(x_1-x_2,t)\frac{d\Lambda}{\Lambda}\right] \\ &+2g\pi I_C(x_1,x_2,t)\frac{d\Lambda}{\Lambda}R_<^{0}(x_1-x_2,t) \end{aligned} \tag{142}$$

Rewriting the zero-order term for convenience,

$$R^{(0)}(x_1-x_2,t)= R_<^{(0)}(x_1-x_2,t)\left[1-\frac{d\Lambda}{\Lambda}\gamma_{an,0}(x_1-x_2,t)\right] \tag{143}$$

and combining Eq. 142, 143, we get

$$\begin{aligned} R =& R_<^{(0)}\left[1-\frac{d\Lambda}{\Lambda}\gamma_{an,0}(x_1-x_2,t)+\frac{d\Lambda}{\Lambda}2g\pi I_C(x_1-x_2,t)\right] \\ &+R_<^{(1)}\left[1+(2-K_{neq})\frac{d\Lambda}{\Lambda}+\left\{-K_{neq}+\gamma_{an,0}(x_1-x_2,t)\right\}\frac{d\Lambda}{\Lambda}\right] \end{aligned} \tag{144}$$

The above implies,

$$\left[\frac{\partial}{\partial\ln\Lambda}+\gamma_{an,0}-2\pi g I_C\right]R(x_1,x_2,T_m)=0 \tag{145}$$

Including the β-function derived in a previous section and which appears at $\mathcal{O}(g^2)$, the above can be rewritten as

$$\left[\frac{\partial}{\partial\ln l}+\beta(g_i)\frac{\partial}{\partial g_i}-\gamma_{an,0}+2\pi g I_C\right]R\left[\frac{r\Lambda_0}{l},\frac{\Lambda_0 T_{m0}}{l},g_i(l)\right]=0 \tag{146}$$

where $g_i=g,K,\eta,T_{eff}$, and Λ_0,T_{m0},g_{i0} denote bare values. Moreover I_C is given by

$$\begin{aligned} &I_C(x_1,x_2,t,K_{neq},K_{eq}) = -\left(\frac{1}{2\pi}\right)e^{K_{neq}\ln\sqrt{1+(x_1-x_2)^2}}\int_{-\infty}^{\infty}dx'\int_0^t dt' \\ &\sin\left[\frac{K_{eq}}{2}\left\{\tan^{-1}(t'+x'-x_1)+\tan^{-1}(t'-(x'-x_1))+\tan^{-1}(t'+x'-x_2)+\tan^{-1}(t'-(x'-x_2))\right\}\right] \\ &\times e^{-\frac{K_{neq}}{2}\left[\ln\sqrt{1+(t'+x'-x_1)^2}+\ln\sqrt{1+(t'-(x'-x_1))^2}+\ln\sqrt{1+(t'+x'-x_2)^2}+\ln\sqrt{1+(t'-(x'-x_2))^2}\right]} \\ &\times e^{-\frac{K_{tr}}{2}\left[\ln\sqrt{\frac{1+(2t-t'+x'-x_1)^2}{1+4t^2}}+\ln\sqrt{\frac{1+(2t-t'-(x'-x_1))^2}{1+4(t-t')^2}}+\ln\sqrt{\frac{1+(2t-t'+x'-x_2)^2}{1+4t^2}}+\ln\sqrt{\frac{1+(2t-t'-(x'-x_2))^2}{1+4(t-t')^2}}\right]} \\ &\times\left[\frac{K_{tr}}{1+4t^2}+\frac{K_{tr}}{1+4(t-t')^2}-\frac{K_{neq}}{2}\left\{\frac{1}{1+(t'+x'-x_1)^2}+\frac{1}{1+(t'-(x'-x_1))^2}+\frac{1}{1+(t'+x'-x_2)^2}\right.\right. \\ &\left.+\frac{1}{1+(t'-(x'-x_2))^2}\right\}-\frac{K_{tr}}{2}\left\{\frac{1}{1+(2t-t'+x'-x_1)^2}+\frac{1}{1+(2t-t'-(x'-x_1))^2}+\frac{1}{1+(2t-t'+x'-x_2)^2}\right. \\ &\left.\left.+\frac{1}{1+(2t-t'-(x'-x_2))^2}\right\}\right] \end{aligned} \tag{147}$$

$$\begin{aligned} &+\frac{1}{2\pi}e^{K_{neq}\ln\sqrt{1+(x_1-x_2)^2}}\int_{-\infty}^{\infty}dx'\int_0^t dt'\cos\left[\frac{K_{eq}}{2}\left\{\tan^{-1}(t'+x'-x_1)+\tan^{-1}(t'-(x'-x_1))\right.\right. \\ &\left.\left.+\tan^{-1}(t'+x'-x_2)+\tan^{-1}(t'-(x'-x_2))\right\}\right] \\ &\times e^{-\frac{K_{neq}}{2}\left[\ln\sqrt{1+(t'+x'-x_1)^2}+\ln\sqrt{1+(t'-(x'-x_1))^2}+\ln\sqrt{1+(t'+x'-x_2)^2}+\ln\sqrt{1+(t'-(x'-x_2))^2}\right]} \\ &\times e^{-\frac{K_{tr}}{2}\left[\ln\sqrt{\frac{1+(2t-t'+x'-x_1)^2}{1+4t^2}}+\ln\sqrt{\frac{1+(2t-t'-(x'-x_1))^2}{1+4(t-t')^2}}+\ln\sqrt{\frac{1+(2t-t'+x'-x_2)^2}{1+4t^2}}+\ln\sqrt{\frac{1+(2t-t'-(x'-x_2))^2}{1+4(t-t')^2}}\right]} \\ &\times\frac{K_{eq}}{2}\left[\frac{(t'+x'-x_1)}{1+(t'+x'-x_1)^2}+\frac{(t'-(x'-x_1))}{1+(t'-(x'-x_1))^2}+\frac{(t'+x'-x_2)}{1+(t'+x'-x_2)^2}+\frac{(t'-(x'-x_2))}{1+(t'-(x'-x_2))^2}\right] \end{aligned} \tag{148}$$

The usual CS equation encountered in equilibrium for the Heisenberg chain [25–27] may be obtained from the above by setting the time after the quench $t=\infty$, and by setting $K_0=K$ and by also in addition being at the critical point $K_{eq}=2$. Here, noting that $-\text{Im}\left[e^{-i\tan^{-1}x}\frac{1}{1+ix}\right]=\frac{\sin\left(\tan^{-1}x\right)}{1+x^2}+\frac{x}{1+x^2}\cos\left(\tan^{-1}x\right)$, one may write

$$I_C=-\frac{\left(1+(x_1-x_2)^2\right)}{2\pi}\text{Im}\left[\int_{-\infty}^{\infty}dx'\int_0^{\infty}dt'e^{-\ln\left(1+i(t'+x'-x_1)\right)-\ln\left(1+i(t'-x'+x_1)\right)-\ln\left(1+i(t'+x'-x_2)\right)-\ln\left(1+i(t'-x'+x_2)\right)}\right.$$
$$\left.\left\{\frac{1}{1+i(t'+x'-x_1)}+\frac{1}{1+i(t'-x'+x_1)}+\frac{1}{1+i(t'+x'-x_2)}+\frac{1}{1+i(t'-x'+x_2)}\right\}\right] \tag{149}$$

Analytically continuing in time,

$$I_C=\frac{1+(x_1-x_2)^2}{2\pi}\int_{-\infty}^{\infty}dx'\int_0^{\infty}d\tau\left[-\frac{\partial}{\partial\tau}\right]\left[\frac{1}{(1+\tau)^2+(x'-x_1)^2}\times\frac{1}{(1+\tau)^2+(x'-x_2)^2}\right] \tag{150}$$

$$=\frac{(x_1-x_2)^2+1}{4+(x_1-x_2)^2} \tag{151}$$

Thus for $|x_1-x_2|\gg 1$,

$$I_C\left(K_0=K,K_{eq}=2,t=\infty\right)=1 \tag{152}$$

Correlation function for the pure lattice quench

We want to solve

$$\left[\frac{\partial}{\partial\ln l}+\beta(g_i)\frac{\partial}{\partial g_i}-\gamma_{an,0}+2\pi gI_C\right]R\left[\frac{(x_1-x_2)\Lambda_0}{l},\frac{\Lambda_0T_{m0}}{l},g_i(l)\right] \tag{153}$$

where $g_i=g,K,\eta,T_{eff}$. In addition, Λ_0,T_{m0},g_{i0} denote bare values. In this section, we want to solve Eq. 153 for the lattice quench $K_0=K$ and near the critical point $K_{eq}=2+\delta$. Note that,

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\ll 1)=\frac{1}{2}\left(K_{neq}+K_{tr}\right)=\frac{\gamma^2K_0}{8}=1+\frac{\delta}{2} \tag{154}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t\gg|x_1-x_2|)=\frac{K_{neq}}{2}=1+\frac{\delta}{2} \tag{155}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t\ll|x_1-x_2|)=\frac{K_{neq}}{2}=1+\frac{\delta}{2} \tag{156}$$

$$\gamma_{an,0}(|x_1-x_2|\Lambda\gg 1,t\Lambda\gg 1,2t=|x_1-x_2|)=\frac{1}{2}\left(K_{neq}-\frac{K_{tr}}{2}\right)=1+\frac{\delta}{2} \tag{157}$$

and for I_C, the full expression in Eqns 147, 148 reduces to

$$I_C(x_1-x_2,t)=\frac{(x_1-x_2)^2+1}{4+(x_1-x_2)^2}-\frac{(x_1-x_2)^2+1}{2\pi}\text{Im}\left[i\int_{-\infty}^{\infty}dx\left\{\frac{1}{(1+it)^2+x^2}\times\frac{1}{(1+it)^2+(x-(x_1-x_2))^2}\right\}\right]$$
$$=\frac{(x_1-x_2)^2+1}{4+(x_1-x_2)^2}-\frac{1+(x_1-x_2)^2}{1+t^2}\left[\frac{(x_1-x_2)^2-4t^2+4-8t^2}{\left\{(x_1-x_2)^2-4t^2+4\right\}^2+64t^2}\right] \tag{158}$$

The following different limits exist,

$$I_C(|x_1-x_2|\gg 1,t\gg 1,t\gg|x_1-x_2|)=1+\mathcal{O}\left(\frac{1}{(x_1-x_2)^2},\frac{(x_1-x_2)^2}{t^4}\right) \tag{159}$$

$$I_C(|x_1-x_2|\gg 1,t\gg 1,t\ll|x_1-x_2|)=1+\mathcal{O}\left(\frac{1}{t^2}\right) \tag{160}$$

$$I_C(|x_1-x_2|\gg 1,t\gg 1,2t=|x_1-x_2|)=1+1/2=3/2+\mathcal{O}\left(\frac{1}{(x_1-x_2)^2}\right) \tag{161}$$

In addition the β-function for a pure lattice quench and using $K_{neq} = K_{eq} = 2 + \delta$ is,

$$\frac{dg}{d\ln l} = -g\delta \tag{162}$$

$$\frac{d\delta}{d\ln l} = -4\pi g^2 I_K(T_m) \tag{163}$$

$$\frac{1}{u}\frac{du}{d\ln l} = 2\pi g^2 I_u(T_m) \tag{164}$$

$$\frac{d\eta}{d\ln l} = \eta + 4\pi g^2 I_\eta(T_m) \tag{165}$$

$$\frac{d(\eta T_{eff})}{d\ln l} = 2\eta T_{eff} + 2\pi g^2 I_{T_{eff}}(T_m) \tag{166}$$

$$\frac{dT_m}{d\ln l} = -T_m \tag{167}$$

where

$$I_K(T_m, K_{neq} = K_{eq} = 2) = \pi - \frac{\pi}{2}\left[\frac{2 + 20T_m^2 + 7 \times 16T_m^4}{(1 + 4T_m^2)^3}\right] \tag{168}$$

thus

$$I_K(T_m \ll 1) = 2\pi T_m^2 + \ldots \tag{169}$$

$$I_K(T_m \gg 1) = \pi\left[1 - \frac{7}{8T_m^2} + \ldots\right] \tag{170}$$

Moreover,

$$I_\eta = 4\pi \frac{(2T_m)^3}{\left[1 + (2T_m)^2\right]^3} \tag{171}$$

$$I_{T_{eff}} = 6\pi T_m \left[\frac{1 - \frac{4}{3}T_m^2}{(1 + 4T_m^2)^3}\right] \tag{172}$$

At very long times, for the pure lattice quench, we may neglect T_{eff}, η as these are zero at steady-state at $\mathcal{O}(g^2)$ [9]. Now, Eq. 153 implies,

$$R\left(\frac{r\Lambda_0}{l}, \frac{T_{m0}\Lambda_0}{l}, g(l)\right) = e^{\int_{g_0}^{g(l)} dg' \frac{\gamma_{an}(g')}{\beta(g')}} R\left(\Lambda_0 r, T_{m0}\Lambda_0, g_0\right) \tag{173}$$

where

$$\gamma_{an} = \gamma_{an,0} - 2\pi g I_C \tag{174}$$

Eq. 173 implies

$$R\left(\Lambda_0 r, \Lambda_0 T_{m0}, g_0\right) = e^{-\int_{g_0}^{g(l)} dg' \frac{\gamma_{an}(g')}{\beta(g')}} R\left(\frac{r\Lambda_0}{l}, \frac{T_{m0}\Lambda_0}{l}, g(l)\right) \tag{175}$$

One interesting case is the behavior of correlators at steady-state $T_{m0} \to \infty$. Here the explicit dependence on the time after the quench drops out. Then integrating upto $l = \Lambda_0 r$, we get

$$R\left(\Lambda_0 r, T_{m0} = \infty, g_0\right) = e^{-\int_{g_0}^{g(\Lambda_0 r)} dg' \frac{\gamma_{an}(g')}{\beta(g')}} R\left(\frac{r\Lambda_0}{l} = 1, T_{m0} = \infty, g(l)\right) \tag{176}$$

where for $l \gg 1$, $g(l)$ being negligibly small, the correlator on the r.h.s. may be evaluated within perturbation theory. Moreover, $R\left(\frac{r\Lambda_0}{l} = 1, T_{m0} = \infty, g(l)\right) = \mathcal{O}(1) + \mathcal{O}\left(g(l)\right) + \ldots$ as this is a short-distance correlator.

Another interesting case is for spatial separations on the light-cone where $r = 2T_{m0}$. Here,

$$\begin{aligned} R\left(\Lambda_0 r, 2T_{m0} = r, g_0\right) &= e^{-\int_{g_0}^{g(\Lambda_0 r)} dg' \frac{\gamma_{an}(g')}{\beta(g')}} \\ &\times R\left(\frac{r\Lambda_0}{l} = 1, 2T_{m0} = l, g(l)\right) \end{aligned} \tag{177}$$

Here R on the r.h.s may again be evaluated within perturbation theory, and being a short-distance correlator, $R = \mathcal{O}(1)$.

At the critical point $\delta = 2\pi g$, so that

$$\frac{d\delta}{d\ln l} = -\delta^2 \tag{178}$$

and

$$\gamma_{an} = 1 + \frac{\delta}{2} - \delta I_C \tag{179}$$

where $I_C = 1$ for $T_{m0} \gg r$, whereas $I_C = 3/2$ for points on the light cone $r = 2T_{m0}$. Thus in general,

$$\frac{\gamma_{an}}{\beta} = -\frac{1}{\delta^2} + \frac{1}{\delta}\left(I_C - \frac{1}{2}\right) \tag{180}$$

The solution of Eq. 178 is

$$\frac{1}{\delta(\Lambda_0 r)} - \frac{1}{\delta_0} = \ln(\Lambda_0 r) \tag{181}$$

while

$$\int_{\delta_0}^{\delta(\Lambda_0 r)} d\delta \frac{\gamma_{an}}{\beta} = \ln(\Lambda_0 r) + \left(I_C - \frac{1}{2}\right) \ln\left[\frac{\delta(\Lambda_0 r)}{\delta_0}\right] \tag{182}$$

Thus the correlation function is found to be

$$R(\Lambda_0 r, g_0) \simeq \frac{\left[\ln(\Lambda_0 r)\right]^{I_C - 1/2}}{\Lambda_0 r} \tag{183}$$

Eq. 183 shows that far from the light cones where $T_{m0} \gg r$, the correlator decays as $\frac{\sqrt{\ln r}}{r}$ as in equilibrium, while right at the light-cone $r = 2T_{m0}$, the correlator decays somewhat slower, as $\frac{\ln r}{r}$.

Correlation function for the lattice and interaction quench

Here we are interested in the regime where $K_{neq} \simeq 2$, which implies

$$K = K_0\sqrt{\frac{16}{\gamma^2 K_0} - 1} \equiv K_{eq} = \frac{\gamma^2 K_0}{4}\sqrt{\frac{16}{\gamma^2 K_0} - 1} \tag{184}$$

$$K_{tr} = \frac{\gamma^2 K_0}{4} - 2 \tag{185}$$

Writing $K_{neq} = 2 + \delta$, the RG equations for times $T_m < 1/\eta$ (*i.e.*, on neglecting dissipative effects) are the following

$$\frac{dg}{d\ln l} = -g\delta \tag{186}$$

$$\frac{d\delta}{d\ln l} = -\pi g^2 \left(\frac{\gamma^2 K_0}{4}\right)^2 \left(\frac{16}{\gamma^2 K_0} - 1\right)^{3/2} I_K(T_m) \tag{187}$$

$$\frac{dT_m}{d\ln l} = -T_m \tag{188}$$

above we have used that since $K_{neq} = 2 + \delta$, $dK = \frac{4}{\gamma^2}\frac{K_0}{K}d\delta$, and $K_{eq} = \frac{\gamma^2 K_0}{4}\sqrt{\frac{16}{\gamma^2 K_0} - 1}$. Moreover,

$$I_K(T_m = \infty, K_{neq} = 2) = -4\int_0^\infty du \frac{u}{(1+u^2)^2}\cos\left(K_{eq}\tan^{-1} u\right)\underbrace{\int_0^u dv \frac{v}{1+v^2}\sin\left(K_{eq}\tan^{-1} v\right)}_{i_1(u)} \tag{189}$$

$$-4\int_0^\infty du \frac{u}{1+u^2}\cos\left(K_{eq}\tan^{-1} u\right)\underbrace{\int_0^u dv \frac{v}{(1+v^2)^2}\sin\left(K_{eq}\tan^{-1} v\right)}_{i_2(u)} \tag{190}$$

$$-2K_{eq}\int_0^\infty du \frac{u}{1+u^2}\cos\left(K_{eq}\tan^{-1} u\right)\underbrace{\int_0^u dv \frac{v^2}{(1+v^2)^2}\cos\left(K_{eq}\tan^{-1} v\right)}_{i_3(u)} \tag{191}$$

$$+2K_{eq}\int_0^\infty du \frac{u^2}{(1+u^2)^2}\sin\left(K_{eq}\tan^{-1} u\right)\underbrace{\int_0^u dv \frac{v}{1+v^2}\sin\left(K_{eq}\tan^{-1} v\right)}_{i_1(u)} \tag{192}$$

Note that Eqns 190, 191 are logarithmically divergent (such a divergence does not arise for a pure lattice quench). The reason for this divergence is because we have used leading order perturbation theory to evaluate the correlators that go into I_K, whereas, we should have used the correlators from the full theory which correspond to a non-zero dissipation η. Taking this into account, the logarthmic divergence is cut-off by $\min(T_m, 1/\eta)$. Denoting the well behaved terms in Eqns 189, 192 as πc_1, and using $i_2(\infty) = -\frac{1}{K_{eq}^2 - 4}\sin\left(\frac{\pi K_{eq}}{2}\right)$ and $i_3(\infty) = \frac{(K_{eq}^2-2)}{K_{eq}(K_{eq}^2-4)}\sin\left(\frac{\pi K_{eq}}{2}\right)$, we may write,

$$I_K(T_m, K_{neq} = 2) = \pi\left[c_1 - 2c_2\sin\left(\frac{\pi K_{eq}}{2}\right)\int_0^{min(T_m,1/\eta)} du\frac{u}{1+u^2}\cos\left(K_{eq}\tan^{-1}u\right)\right] \tag{193}$$

$$= \pi\left[c_1 - c_2\sin\left(\pi K_{eq}\right)\ln\left(\min(T_m, 1/\eta)\right)\right] \tag{194}$$

where $c_{1,2}$ are $\mathcal{O}(1)$ and depend on K_{eq} and hence K_0. When $K_0 = K$, $c_1 = 1$.

In what follows we will consider a prethermalized regime where $T_m < 1/\eta$, and $|K_{eq} - K_{neq}|\ln T_m \ll 1$. In this case, $I_K \simeq \pi c_1$. The solution of the RG equations in this regime, under conditions that

$$g_{eff} = gB < \delta \tag{195}$$

where

$$B = \pi\sqrt{c_1}\left(\frac{\gamma^2 K_0}{4}\right)\left(\frac{16}{\gamma^2 K_0} - 1\right)^{3/4} \tag{196}$$

and defining a constant of the RG flow $\frac{dg_{eff}}{d\ln l} = -g_{eff}\delta$, $\frac{d\delta}{d\ln l} = -g_{eff}^2$,

$$A = \sqrt{\delta^2 - g_{eff}^2} \tag{197}$$

is given by,

$$\delta(l) = A\coth\left[A\ln l + \tanh^{-1}\left(\frac{A}{\delta_0}\right)\right] \tag{198}$$

$$g_{eff}(l) = \frac{A}{\sinh\left[A\ln l + \tanh^{-1}\left(\frac{A}{\delta_0}\right)\right]} \tag{199}$$

We will now study the system in the vicinity of the non-equilibrium critical point $K_{neq} = 2 + \delta$. Here for I_C we find

$$I_C(r \gg 1, T_m \gg 1, T_m \gg r) = \frac{K_{eq}}{2} + \left[1 - \left(\frac{K_{eq}}{2}\right)^2\right]L(K_{eq}) \tag{200}$$

where

$$L(K_{eq}) = \lim_{|r|\gg 1}\left(\frac{r^2}{4\pi}\int_0^\infty du\int_{-u}^{u} dv\,\sin\left[\frac{K_{eq}}{2}\left\{\tan^{-1}u+\tan^{-1}v+\tan^{-1}(u+r)+\tan^{-1}(v-r)\right\}\right]\right.$$
$$\left.\frac{1}{\sqrt{1+u^2}}\frac{1}{\sqrt{1+v^2}}\frac{1}{\sqrt{1+(u+r)^2}}\frac{1}{\sqrt{1+(v-r)^2}}\left[\frac{1}{1+u^2}+\frac{1}{1+v^2}+\frac{1}{1+(u+r)^2}+\frac{1}{1+(v-r)^2}\right]\right) \tag{201}$$
$$+(r\to -r)$$

$L(K_{eq})$ is a universal function of K_{eq} and hence the initial Luttinger parameter. I_C is plotted in the main text.

The solution of the CZ equations is,

$$R\left(\Lambda_0 r, \Lambda_0 T_{m0}, g_0\right) = e^{-\int_{g_0}^{g(l)} dg'\frac{\gamma_{an}(g')}{\beta(g')}} R\left(\frac{r\Lambda_0}{l}, \frac{T_{m0}\Lambda_0}{l}, g(l)\right) \tag{202}$$

We will solve this for $1/\eta \gg T_{m0} \gg r$. For this case, dissipative effects may be neglected in the solution of the RG equations which will be integrated upto $l=\Lambda_0 r$. In particular,

$$R\left(\Lambda_0 r, r\ll T_{m0}\ll 1/\eta, g_0\right) = e^{-\int_{g_0}^{g(\Lambda_0 r)} dg'\frac{\gamma_{an}(g')}{\beta(g')}} R\left(\frac{r\Lambda_0}{l}=1, \frac{T_{m0}}{r}\gg 1, g(l)\right) \tag{203}$$

In the r.h.s, R being a short-distance correlator, is $\mathcal{O}(1)$. Note that

$$\gamma_{an}(r\ll T_{m0}) = 1+\frac{\delta}{2}-2\pi g I_C \tag{204}$$

Then,

$$\int_{g_0}^{g(l)}\frac{\gamma(g')}{\beta(g')}dg' = \ln l + \frac{1}{2}\ln\left[\frac{\sinh\left(A\ln l+\tanh^{-1}\frac{A}{\delta_0}\right)}{\sinh\left(\tanh^{-1}\frac{A}{\delta_0}\right)}\right] - \frac{2\pi I_C A}{B}\ln\left[\frac{\tanh\left(\frac{A}{2}\ln l+\frac{1}{2}\tanh^{-1}\frac{A}{\delta_0}\right)}{\tanh\left(\frac{1}{2}\tanh^{-1}\frac{A}{\delta_0}\right)}\right] \tag{205}$$

where $B=\pi\sqrt{c_1}\left(\frac{\gamma^2K_0}{4}\right)\left(\frac{16}{\gamma^2K_0}-1\right)^{3/4}$.

The correlation function within the light-cone is given by setting $l=r$,

$$R\left(\Lambda_0 r, r\ll T_{m0}\ll 1/\eta, g_0\right) \simeq \frac{1}{(\Lambda_0 r)^{1+A/2}}\sqrt{\frac{1}{1-(\Lambda_0 r)^{-2A}}}\left[\frac{1-(\Lambda_0 r)^{-A}}{1+(\Lambda_0 r)^{-A}}\right]^{\frac{2\pi I_C}{B}} \tag{206}$$

For $A\ln r\ll 1$, we obtain the following logarithmic correction to scaling,

$$R\sim\frac{1}{r}\left(\ln r\right)^{\frac{2\pi I_c}{B}-\frac{1}{2}} \tag{207}$$

Let us now discuss how I_C behaves for spatial separations on the light-cone. Simplifying the full expression in Eqns. 147, 148 by noting that for $r\gg 1, t\gg 1, r=2t$,

$$\sqrt{\frac{1+\left(2t-t'+x'\right)^2}{1+4t^2}}\simeq 1$$
$$\sqrt{\frac{1+\left(2t-t'-x'\right)^2}{1+4(t-t')^2}}\simeq 1$$
$$\sqrt{\frac{1+\left(2t-t'+x'-r\right)^2}{1+4t^2}} = \sqrt{\frac{1+\left(t'-x'\right)^2}{1+4t^2}}$$
$$\sqrt{\frac{1+\left(2t-t'-x'+r\right)^2}{1+4(t-t')^2}}\simeq 2$$

one may write,

$$
\begin{aligned}
& I_C(r=2t, r \gg 1, K_{neq}=2) = 2^{-K_{tr}/2} \mathrm{Lt}_{r\to\infty} \frac{r^{2+\frac{K_{tr}}{2}}}{2\pi} \int_{-\infty}^{\infty} dx' \int_0^{r/2} dt' \\
& \sin\left[\frac{K_{eq}}{2}\left\{\tan^{-1}(t'+x') + \tan^{-1}(t'-x') + \tan^{-1}(t'+x'-r) + \tan^{-1}(t'-x'+r)\right\}\right] \\
& \times \frac{1}{\sqrt{1+(t'+x')^2}} \left(\frac{1}{\sqrt{1+(t'-x')^2}}\right)^{1+\frac{K_{tr}}{2}} \frac{1}{\sqrt{1+(t'+x'-r)^2}} \frac{1}{\sqrt{1+(t'-x'+r)^2}} \\
& \times \left\{\frac{1}{1+(t'+x')^2} + \frac{\left(1+\frac{K_{tr}}{2}\right)}{1+(t'-x')^2} + \frac{1}{1+(t'+x'-r)^2} + \frac{1}{1+(t'-x'+r)^2}\right\} \\
& + 2^{-K_{tr}/2} \mathrm{Lt}_{r\to\infty} \frac{r^{2+\frac{K_{tr}}{2}}}{2\pi} \left(\frac{K_{eq}}{2}\right) \int_{-\infty}^{\infty} dx' \int_0^{r/2} dt' \\
& \cos\left[\frac{K_{eq}}{2}\left\{\tan^{-1}(t'+x') + \tan^{-1}(t'-x') + \tan^{-1}(t'+x'-r) + \tan^{-1}(t'-x'+r)\right\}\right] \\
& \times \frac{1}{\sqrt{1+(t'+x')^2}} \left(\frac{1}{\sqrt{1+(t'-x')^2}}\right)^{1+\frac{K_{tr}}{2}} \frac{1}{\sqrt{1+(t'+x'-r)^2}} \frac{1}{\sqrt{1+(t'-x'+r)^2}} \\
& \times \left[\frac{(t'+x')}{1+(t'+x')^2} + \frac{(t'-x')}{1+(t'-x')^2} + \frac{(t'+x'-r)}{1+(t'+x'-r)^2} + \frac{(t'-x'+r))}{1+(t'-x'+r)^2}\right]
\end{aligned}
\tag{208}
$$

Here we find that

$$
I_C(r \gg 1) \sim r^{K_{tr}/2} \tag{209}
$$

Thus the scaling for spatial separations on the light-cone $r = 2t$ which we found for a pure lattice quench, is lost for a simultaneous lattice and interaction quench when $K_{tr} > 0$ (i.e., $\frac{\gamma^2 K_0}{4} > 2$ so that the initial state is irrelevant). In contrast, scaling continues to hold for spatial separations inside the light-cone.